arXiv:1612.03295v1 [math.AP] 10 Dec 2016

Local Uniqueness and Refined Spike Profiles of Ground States for Two-Dimensional Attractive Bose-Einstein Condensates

Yujin Guo*, Changshou Lin†, and Juncheng Wei‡

Abstract

We consider ground states of two-dimensional Bose-Einstein condensates with attractive interactions, which can be described equivalently by positive minimizers of the L^2-critical constraint Gross-Pitaevskii energy functional. The local uniqueness of ground states is proved as $a \nearrow a^* := \|w\|_2^2$ for a class of homogeneous trapping potentials, where a denotes the interaction strength and w is the unique positive solution of $\Delta w - w + w^3 = 0$ in $\mathbb{R}^2$. The refined spike profiles of ground states as $a \nearrow a^*$ are also derived rigorously under more general trapping potentials.

Keywords: Bose-Einstein condensation; spike profiles; local uniqueness; Pohozaev identity.

1 Introduction

The phenomenon of Bose-Einstein condensation (BEC) has been investigated intensively since its first realization in cold atomic gases, see [1, 5] and references therein. In these experiments, a large number of (bosonic) atoms are confined to a trap and cooled to very low temperatures. Condensation of a large fraction of particles into the same one-particle state is observed below a critical temperature. These Bose-Einstein condensates display various interesting quantum phenomena, such as the critical-mass collapse, the superfluidity and the appearance of quantized vortices in rotating traps (e.g.[5]). Specially, if the force between the atoms in the condensates is attractive, the system collapses as soon as the particle number increases beyond a critical value, see, e.g., [23] or [5, Sec. III.B].

Bose-Einstein condensates (BEC) with attractive interactions in $\mathbb{R}^2$ can be described ([2, 5, 10]) by the following Gross-Pitaevskii (GP) energy functional

$$E_a(u) := \int_{\mathbb{R}^2} \Big(|\nabla u|^2 + V(x)|u|^2 \Big) dx - \frac{a}{2} \int_{\mathbb{R}^2} |u|^4 dx, \tag{1.1}$$

*Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, P.O. Box 71010, Wuhan 430071, P. R. China. Email: yjguo@wipm.ac.cn. Y. J. Guo is partially supported by NSFC grants 11322104 and 11671394.

†Taida Institute of Mathematical Sciences, National Taiwan University, Taipei 10617, Taiwan. Email: cslin@math.ntu.edu.tw.

‡Department of Mathematics, University of British Columbia, Vancouver, BC V6T 1Z2, Canada. Email: jcwei@math.ubc.ca. J.C. Wei is partially supported by NSERC of Canada.

where $a > 0$ describes the strength of the attractive interactions, and $V(x)$ denotes the trapping potential satisfying $\lim_{|x|\to\infty} V(x) = \infty$. As addressed recently in [10, 11], ground states of attractive BEC in $\mathbb{R}^2$ can be described by the constraint minimizers of the GP energy

$$e(a) := \inf_{\{u\in\mathcal{H},\, \|u\|_2^2=1\}} E_a(u), \tag{1.2}$$

where the space $\mathcal{H}$ is defined by

$$\mathcal{H} := \Big\{u \in H^1(\mathbb{R}^2) : \int_{\mathbb{R}^2} V(x)|u(x)|^2 dx < \infty\Big\}. \tag{1.3}$$

The minimization problem $e(a)$ was analyzed recently in [2, 10, 11, 12, 26] and references therein. Existing results show that $e(a)$ is an L^2-critical constraint variational problem. Actually, it was shown in [2, 10] that $e(a)$ admits minimizers if and only if $a < a^* := \|w\|_2^2$, where $w = w(|x|)$ is the unique (up to translations) radial positive solution (cf. [7, 19, 14]) of the following nonlinear scalar field equation

$$\Delta w - w + w^3 = 0 \ \text{ in } \ \mathbb{R}^2, \ \text{ where } \ w \in H^1(\mathbb{R}^2)\,. \tag{1.4}$$

It turns out that the existence and nonexistence of minimizers for $e(a)$ are well connected with the following Gagliardo-Nirenberg inequality

$$\int_{\mathbb{R}^2} |u(x)|^4 dx \le \frac{2}{\|w\|_2^2} \int_{\mathbb{R}^2} |\nabla u(x)|^2 dx \int_{\mathbb{R}^2} |u(x)|^2 dx, \ \ u \in H^1(\mathbb{R}^2)\,, \tag{1.5}$$

where the equality is attained at w (cf. [25]).

Since $E_a(u) \ge E(|u|)$ for any $u \in \mathcal{H}$, without loss of generality one can restrict the minimizations of $e(a)$ to positive functions in view of the maximum principle. Therefore, in this paper positive minimizers of $e(a)$ are called ground states of attractive BEC. If u_a is a positive minimizer of $e(a)$ for $a < a^*$, it then satisfies the Euler-Lagrange equation

$$-\Delta u_a + V(x)u_a = \mu_a u_a + a u_a^3 \quad \text{ in } \ \mathbb{R}^2, \tag{1.6}$$

where $\mu_a \in \mathbb{R}$ is a suitable Lagrange multiplier. Applying energy estimates and blow-up analysis, the spike profiles of positive minimizers for $e(a)$ as $a \nearrow a^*$ were recently discussed in [10, 11, 12] under different types of potentials $V(x)$, see our Proposition 2.2 for some related results. In spite of these facts, it remains open to discuss the refined spike profiles of positive minimizers. On the other hand, the local uniqueness of positive minimizers for $e(a)$ as *a.e.* $a \nearrow a^*$ was also proved [11] by the ODE argument, for the case where $V(r) = V(|x|)$ is radially symmetric and satisfies $V'(r) \ge 0$, see Corollary 1.1 in [11] for details. It is therefore natural to ask whether such local uniqueness still holds for the case where $V(x)$ is not radially symmetric. We should remark that all these results mentioned above were obtained mainly by analyzing the variational structures of the minimization problem $e(a)$, instead of discussing the PDE properties of the associated elliptic equation (1.6).

By investigating thoroughly the associated equation (1.6), the main purpose of this paper is to derive the refined spike profiles of positive minimizers for $e(a)$ as $a \nearrow a^*$, and extend the above local uniqueness to the cases of non-radial potentials $V(x)$ as well. We shall consider the trapping potential $V(x)$ in the class of homogeneous functions, for which we define

Definition 1.1. $h(x) \geq 0$ in $\mathbb{R}^2$ is homogeneous of degree $p \in \mathbb{R}^+$ about the origin, if there exists some $p > 0$ such that

$$h(tx) = t^p h(x) \text{ in } \mathbb{R}^2 \text{ for any } t > 0. \tag{1.7}$$

The above definition implies that the homogeneous function $h(x)$ of degree $p > 0$ satisfies

$$0 \leq h(x) \leq C|x|^p \text{ in } \mathbb{R}^2, \tag{1.8}$$

where $C > 0$ denotes the maximum of $h(x)$ on $\partial B_1(0)$, and $\nabla h(x) = 0$ if and only if $x = 0$ for the case where $\lim_{|x|\to\infty} h(x) = \infty$.

To introduce our main results, for convenience we denote

$$\lambda_0 = \left(\frac{p}{2}\int_{\mathbb{R}^2} h(x)w^2(x)dx\right)^{\frac{1}{2+p}}, \tag{1.9}$$

and

$$\psi(x) = \varphi(x) - \frac{C^*}{2}\big[w(x) + x\cdot\nabla w(x)\big],$$

where $\varphi(x)$ is the unique solution of

$$\nabla\varphi(0) = 0 \text{ and } \big[-\Delta + (1-3w^2)\big]\varphi(x) = -\frac{2w^3}{\int_{\mathbb{R}^2} w^4} - \frac{h(x)w}{\int_{\mathbb{R}^2} h(x)w^2} \text{ in } \mathbb{R}^2, \tag{1.10}$$

and the nonzero constant C^* is given by

$$C^* = \frac{2}{2+p}\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\varphi^2\Big)$$

with ψ_3 being the unique solution of (3.29). Using above notations, our main results of the present paper can be stated as the following theorem.

Theorem 1.1. *Suppose $V(x) = h(x) \in C^2_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and satisfies*

$$\det Jac\, H(0) \neq 0, \tag{1.11}$$

where the vector function $H(y)$ is given by

$$H(y) = \Big(\int_{\mathbb{R}^2}\Big[\frac{\partial h(x+y)}{\partial y_j} - \frac{p-1}{p\int_{\mathbb{R}^2} h(x)w^2}h(x+y)\Big]w^2(x)dx\Big)_{j=1,2}. \tag{1.12}$$

Then there exists a unique positive minimizer u_a for $e(a)$ as $a \nearrow a^*$. Moreover, u_a satisfies

$$\begin{aligned} u_a(x) = \frac{\lambda_0}{\|w\|_2}\Big\{&\frac{1}{(a^*-a)^{\frac{1}{2+p}}}w\Big(\frac{\lambda_0(x-x_a)}{(a^*-a)^{\frac{1}{2+p}}}\Big) + (a^*-a)^{\frac{1+p}{2+p}}\psi\Big(\frac{\lambda_0(x-x_a)}{(a^*-a)^{\frac{1}{2+p}}}\Big)\\ &+(a^*-a)^{\frac{3+2p}{2+p}}\phi_0\Big(\frac{\lambda_0(x-x_a)}{(a^*-a)^{\frac{1}{2+p}}}\Big)\Big\} + o\big((a^*-a)^{\frac{3+2p}{2+p}}\big) \quad as \quad a \nearrow a^* \end{aligned} \tag{1.13}$$

uniformly in $\mathbb{R}^2$ for some function ϕ_0, and the sequence $\{x_a\}$ satisfies

$$\frac{x_a}{(a^*-a)^{\frac{3+p}{2+p}}} = \begin{cases} O(y_0), & \text{if } h(x) \text{ is not symmetric about the origin;}\\ o(1), & \text{if } h(x) \text{ is symmetric about the origin} \end{cases} \tag{1.14}$$

as $a \nearrow a^*$ for some $y_0 \in \mathbb{R}^2$.

Remark 1.1. *The following examples illustrate that (1.11) holds for many classes of non-radial potentials $h(x)$: for $x = (x_1, x_2) \in \mathbb{R}^2$,*

(1). if $h(x) = x_1^{2n} + \lambda x_2^{2n}$, where $\lambda > 0$ and $n \in \mathbb{N}$, then

$$\det Jac\, H(0) = 4\lambda n^2(2n-1)^2 \int_{\mathbb{R}^2} x_1^{2(n-1)} w^2(x) \int_{\mathbb{R}^2} x_2^{2(n-1)} w^2(x) > 0;$$

(2). if $h(x) = (x_1 + \lambda x_2)^2 + x_2^2$, where $\lambda \in \mathbb{R}$, then

$$\det Jac\, H(0) = 4\Big(\int_{\mathbb{R}^2} w^2 \Big)^2 > 0;$$

(3). if $h(x) = (x_1^2 + x_2^2)^3 - \frac{1}{2} x_1 (x_1^2 + x_2^2)^{\frac{5}{2}} \geq 0$, then

$$\begin{aligned} \det Jac\, H(0) = 6 \int_{\mathbb{R}^2} (x_1^2 + 5x_2^2)(x_1^2 + x_2^2) w^2 \Big[6 \int_{\mathbb{R}^2} (5x_1^2 + x_2^2)(x_1^2 + x_2^2) w^2 \\ + \frac{5}{12 \int_{\mathbb{R}^2} h(x) w^2} \int_{\mathbb{R}^2} (6x_1^2 + x_2^2)(x_1^2 + x_2^2)^{\frac{3}{2}} w^2 \Big] > 0. \end{aligned}$$

Theorem 1.1 therefore shows that the local uniqueness of positive minimizers for $e(a)$ as $a \nearrow a^*$ holds for many non-radial potentials $V(x)$.

Theorem 1.1 implies that the mass concentration of minimizers results in the local uniqueness of minimizers. Theorem 1.1 follows directly by combining Theorems 2.1, 3.6 and 4.4, where ϕ_0 is given explicitly. The proof of such local uniqueness is motivated by [3, 6, 9]. Roughly speaking, as derived in Proposition 2.2 we shall first obtain some estimates on the spike behavior of positive minimizers. Under the additional assumption (1.11), the local uniqueness is then proved in Subsection 2.1 by establishing various types of local Pohozaev identities. We remark that even though Pohozaev identity was used widely for obtaining the existence, nonexistence and local uniqueness of solutions (see e.g. [3, 6] and references therein), as far as we know, this is the first work to employ it for discussing the local uniqueness of constraint minimizers.

It should be pointed out that the assumption (1.11) is unnecessary for the refined spike profiles (1.13) and (1.14) of positive minimizers. Actually, we are able to derive rigorously such types of refined spike profiles under more general trapping potentials $V(x)$. More precisely, when $V(x) = h(x)$ is homogeneous of degree p and is non-symmetric about the origin, we shall obtain such refined spike profiles in Section 3, see Theorem 3.6 for more details. However, when the homogeneous function $h(x)$ is symmetric about the origin, as stated by Theorem 4.4, we shall be able to prove such results under the perturbation of $h(x)$, *i.e.*, $V(x) = g(x)h(x)$ satisfying (V) defined in Section 4. To establish Theorem 3.6 and Theorem 4.4, which then imply directly (1.13) and (1.14), our Proposition 2.2 shows that the arguments of [10, 11, 12] give the leading expansion term of the minimizer u_a and the associated Lagrange multiplier μ_a satisfying (1.6) as well. In order to get (1.13) for the rest terms of u_a, the difficulty is to obtain the more precise estimate of μ_a, which is overcome by the very delicate analysis of the associated equation (1.6), together with the constraint condition of u_a.

This paper is organized as follows: In Section 2 we shall prove Theorem 2.1, *i.e.*, the local uniqueness of Theorem 1.1, under the additional assumption (1.11). Section 3 is concerned with proving Theorem 3.6 on the refined spike profiles of positive minimizers for $e(a)$ as $a \nearrow a^*$, for the case where $V(x) = h(x)$ is homogeneous of degree p and is non-symmetric about the origin. The main aim of Section 4 is to derive Theorem 4.4 on the refined spike profiles of positive minimizers under the assumption (V) defined in Section 4. We shall leave the proof of Lemma 3.4 to Appendix A.

2 Local Uniqueness of Positive Minimizers

This section is concerned with the following local uniqueness of positive minimizers for $e(a)$ as $a \nearrow a^*$, provided that $V(x) = h(x)$ is homogeneous and satisfies (1.11). We remark that our argument is applicable definitely to more general types of potentials $V(x)$, which is however beyond the discussion range of the present work.

Theorem 2.1. *Suppose $V(x) = h(x) \in C^2_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and satisfies (1.11). Then there exists a unique positive minimizer u_a for $e(a)$ as $a \nearrow a^*$.*

In order to prove above Theorem 2.1, which then gives the local uniqueness of Theorem 1.1, we need some estimates of positive minimizers for $e(a)$ as $a \nearrow a^*$, which hold essentially for more general potential $V(x) \in C^1_{loc}(\mathbb{R}^2)$ satisfying

$$\begin{aligned} &V(x) = g(x)h(x), \text{ where } 0 < C \le g(x) \le \tfrac{1}{C} \text{ in } \mathbb{R}^2 \text{ and } h(x) \text{ is homogeneous of} \\ &\text{degree } p \in \mathbb{R}^+ \text{ about the origin.} \end{aligned} \tag{2.1}$$

For convenience, we always denote $\{u_k\}$ to be a positive minimizer sequence of $e(a_k)$ with $a_k \nearrow a^*$ as $k \to \infty$, and define

$$\lambda = \left(\frac{pg(0)}{2} \int_{\mathbb{R}^2} h(x) w^2(x) dx \right)^{\frac{1}{2+p}}, \tag{2.2}$$

where $V(x) = g(x)h(x)$ is assumed to satisfy (2.1) with $p > 0$. We also recall from (1.4) that $w(|x|)$ satisfies

$$\int_{\mathbb{R}^2} |\nabla w|^2 dx = \int_{\mathbb{R}^2} |w|^2 dx = \frac{1}{2} \int_{\mathbb{R}^2} |w|^4 dx, \tag{2.3}$$

see also Lemma 8.1.2 in [4]. Moreover, it follows from [7, Prop. 4.1] that w admits the following exponential decay

$$w(x)\,,\ |\nabla w(x)| = O(|x|^{-\frac{1}{2}} e^{-|x|}) \quad \text{as} \quad |x| \to \infty. \tag{2.4}$$

Proposition 2.2. *Suppose $V(x) \in C^1_{loc}(\mathbb{R}^2)$ satisfies (2.1) with $p > 0$, and let $\lambda > 0$ be defined by (2.2). Then there exist a subsequence, still denoted by $\{a_k\}$, of $\{a_k\}$ and $\{x_k\} \subset \mathbb{R}^2$ such that*

(I). The subsequence $\{u_k\}$ satisfies

$$(a^* - a_k)^{\frac{1}{2+p}} u_k \left(x_k + x(a^* - a_k)^{\frac{1}{2+p}} \right) \to \frac{\lambda w(\lambda x)}{\|w\|_2} \ \text{ as } \ k \to \infty \tag{2.5}$$

uniformly in $\mathbb{R}^2$, *and* $\{x_k\}$ *satisfies*

$$\lim_{k\to\infty} \frac{x_k}{(a^*-a_k)^{\frac{1}{2+p}}} = 0. \tag{2.6}$$

Moreover, u_k *satisfies*

$$(a^*-a_k)^{\frac{1}{2+p}} u_k\left(x_k + x(a^*-a_k)^{\frac{1}{2+p}}\right) \le Ce^{-\frac{\lambda}{2}|x|} \ \text{ in } \ \mathbb{R}^2, \tag{2.7}$$

where the constant $C>0$ *is independent of* k.

(II). The energy $e(a_k)$ *satisfies*

$$\lim_{k\to\infty} \frac{e(a_k)}{(a^*-a_k)^{p/(2+p)}} = \frac{\lambda^2}{a^*}\frac{p+2}{p}. \tag{2.8}$$

Proof. Since the proof of Proposition 2.2 is similar to those in [10, 11, 12], which handle (1.1) with different potentials $V(x)$, we shall briefly sketch the structure of the proof.

If $V(x) \in C^1_{loc}(\mathbb{R}^2)$ satisfies (2.1) with $p>0$, we note that $h(x)\ge 0$ satisfies (1.8). Take the test function

$$u_\tau(x) = A_\tau \frac{\tau}{\|w\|_2}\varphi(x)w(\tau x),$$

where the nonnegative cut-off function $\varphi \in C_0^\infty(\mathbb{R}^2)$ satisfies $0\le\varphi(x)\le 1$ in $\mathbb{R}^2$, and $A_\tau>0$ is chosen so that $\int_{\mathbb{R}^2} u_\tau(x)^2dx = 1$. The same proof of Lemma 3 in [10] then yields that

$$e(a) \le C(a^*-a)^{\frac{p}{p+2}} \ \text{ for } \ 0\le a<a^*, \tag{2.9}$$

where the constant $C>0$ is independent of a. By (2.9), we can follow Lemma 4 in [10] to derive that there exists a positive constant K, independent of a, such that

$$\int_{\mathbb{R}^2} |u_a(x)|^4dx \le \frac{1}{K}(a^*-a)^{-\frac{2}{p+2}} \ \text{ for } \ 0\le a<a^*, \tag{2.10}$$

where $u_a>0$ is any minimizer of $e(a)$. Applying (2.9) and (2.10), a proof similar to that of Theorem 2.1 in [12] then gives that there exist two positive constants $m<M$, independent of a, such that

$$m(a^*-a)^{\frac{p}{p+2}} \le e(a) \le M(a^*-a)^{\frac{p}{p+2}} \ \text{ for } \ 0\le a<a^*. \tag{2.11}$$

Based on (2.11), similar to Theorems 1.2 and 1.3 in [12], one can further deduce that there exist a subsequence (still denoted by $\{a_k\}$) of $\{a_k\}$ and $\{x_k\}\subset\mathbb{R}^2$, where $a_k \nearrow a^*$ as $k\to\infty$, such that (2.6)–(2.8) hold, and

$$(a^*-a_k)^{\frac{1}{2+p}} u_k\left(x_k + x(a^*-a_k)^{\frac{1}{2+p}}\right) \to \frac{\lambda w(\lambda x)}{\|w\|_2} \ \text{ strongly in } \ H^1(\mathbb{R}^2) \tag{2.12}$$

as $k\to\infty$. Finally, since w decays exponentially, the standard elliptic regularity theory applied to (2.12) yields that (2.5) holds uniformly in $\mathbb{R}^2$ (e.g. Lemma 4.9 in [18] for similar arguments), and the proof is therefore complete. □

2.1 Proof of local uniqueness

Following Proposition 2.2, this subsection is devoted to the proof of Theorem 2.1. Our proof is stimulated by [3, 6, 9]. We first define the linearized operator $\mathcal{L}$ by

$$\mathcal{L} := -\Delta + (1 - 3w^2) \ \text{ in } \mathbb{R}^2,$$

where $w = w(|x|) > 0$ is the unique positive solution of (1.4) and w satisfies the exponential decay (2.4). Recall from [14, 20] that

$$ker(\mathcal{L}) = span\Big\{\frac{\partial w}{\partial x_1}, \frac{\partial w}{\partial x_2}\Big\}. \tag{2.13}$$

One can note that u_k solves the Euler-Lagrange equation

$$-\Delta u_k(x) + V(x)u_k(x) = \mu_k u_k(x) + a_k u_k^3(x) \ \text{ in } \mathbb{R}^2, \tag{2.14}$$

where $\mu_k \in \mathbb{R}$ is a suitable Lagrange multiplier and satisfies

$$\mu_k = e(a_k) - \frac{a_k}{2}\int_{\mathbb{R}^2} u_k^4(x)dx. \tag{2.15}$$

Moreover, under the more general assumption (2.1) with $p > 0$, one can derive from (2.3) and (2.5) that u_k satisfies

$$\int_{\mathbb{R}^2} u_k^4(x)dx = (a^* - a_k)^{-\frac{2}{2+p}}\Big[\frac{2\lambda^2}{a^*} + o(1)\Big] \ \text{ as } \ k \to \infty. \tag{2.16}$$

It then follows from (2.3), (2.15) and (2.16) that μ_k satisfies

$$\frac{\mu_k \varepsilon_k^2}{\lambda^2} \to -1 \ \text{ as } \ k \to +\infty. \tag{2.17}$$

Set

$$\bar{u}_k(x) := \frac{\sqrt{a^*}\varepsilon_k}{\lambda} u_k\Big(\frac{\varepsilon_k}{\lambda}x + x_k\Big),$$

so that Proposition 2.2 gives $\bar{u}_k(x) \to w(x)$ uniformly in $\mathbb{R}^2$ as $k \to \infty$. Note from (2.14) that $\bar{u}_k$ satisfies

$$-\Delta \bar{u}_k(x) + \Big(\frac{\varepsilon_k}{\lambda}\Big)^2 V\Big(\frac{\varepsilon_k}{\lambda}x + x_k\Big)\bar{u}_k(x) = \frac{\mu_k\varepsilon_k^2}{\lambda^2}\bar{u}_k(x) + \frac{a_k}{a^*}\bar{u}_k^3(x) \ \text{ in } \mathbb{R}^2. \tag{2.18}$$

Moreover, by the exponential decay (2.7), there exist $C_0 > 0$ and $R > 0$ such that

$$|\bar{u}_k(x)| \le C_0 e^{-\frac{|x|}{2}} \ \text{ for } \ |x| > R, \tag{2.19}$$

which then implies that

$$\Big|\Big(\frac{\varepsilon_k}{\lambda}\Big)^2 V\Big(\frac{\varepsilon_k}{\lambda}x + x_k\Big)\bar{u}_k(x)\Big| \le CC_0 e^{-\frac{|x|}{4}} \ \text{ for } \ |x| > R,$$

if $V(x)$ satisfies (2.1) with $p > 0$. Therefore, under the assumption (2.1) with $p > 0$, applying the local elliptic estimates to (2.18) yields that

$$|\nabla \bar{u}_k(x)| \le Ce^{-\frac{|x|}{4}} \ \text{ as } \ |x| \to \infty, \tag{2.20}$$

where the estimates (2.17) and (2.19) are also used. In the following, we shall follow Proposition 2.2 and (2.20) to derive the local uniqueness of positive minimizers as $a \nearrow a^*$.

Proof of Theorem 2.1. Suppose that there exist two different positive minimizers $u_{1,k}$ and $u_{2,k}$ of $e(a_k)$ with $a_k \nearrow a^*$ as $k \to \infty$. Let $x_{1,k}$ and $x_{2,k}$ be the unique local maximum point of $u_{1,k}$ and $u_{2,k}$, respectively. Following (2.14), $u_{i,k}$ then solves the Euler-Lagrange equation

$$-\Delta u_{i,k}(x) + h(x)u_{i,k}(x) = \mu_{i,k}u_{i,k}(x) + a_k u_{i,k}^3(x) \quad \text{in } \mathbb{R}^2, \quad i = 1, 2, \tag{2.21}$$

where $\mu_{i,k} \in \mathbb{R}$ is a suitable Lagrange multiplier. Define

$$\bar{u}_{i,k}(x) := \frac{\sqrt{a^*}\varepsilon_k}{\lambda} u_{i,k}(x), \quad \text{where } i = 1, 2. \tag{2.22}$$

Proposition 2.2 then implies that $\bar{u}_{i,k}\big(\frac{\varepsilon_k}{\lambda}x + x_{2,k}\big) \to w(x)$ uniformly in $\mathbb{R}^2$, and $\bar{u}_{i,k}$ satisfies the equation

$$-\varepsilon_k^2\Delta\bar{u}_{i,k}(x) + \varepsilon_k^2 h(x)\bar{u}_{i,k}(x) = \mu_{i,k}\varepsilon_k^2\bar{u}_{i,k}(x) + \frac{\lambda^2 a_k}{a^*}\bar{u}_{i,k}^3(x) \quad \text{in } \mathbb{R}^2, \quad i = 1, 2. \tag{2.23}$$

Because $u_{1,k} \not\equiv u_{2,k}$, we consider

$$\bar{\xi}_k(x) = \frac{u_{2,k}(x) - u_{1,k}(x)}{\|u_{2,k} - u_{1,k}\|_{L^\infty(\mathbb{R}^2)}} = \frac{\bar{u}_{2,k}(x) - \bar{u}_{1,k}(x)}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}}.$$

Then $\bar{\xi}_k$ satisfies the equation

$$-\varepsilon_k^2\Delta\bar{\xi}_k + \bar{C}_k(x)\bar{\xi}_k = \bar{g}_k(x) \quad \text{in } \mathbb{R}^2, \tag{2.24}$$

where the coefficient $\bar{C}_k(x)$ satisfies

$$\bar{C}_k(x) := -\mu_{1,k}\varepsilon_k^2 - \frac{\lambda^2 a_k}{a^*}\big(\bar{u}_{2,k}^2 + \bar{u}_{2,k}\bar{u}_{1,k} + \bar{u}_{1,k}^2\big) + \varepsilon_k^2 h(x), \tag{2.25}$$

and the nonhomogeneous term $\bar{g}_k(x)$ satisfies

$$\begin{aligned}\bar{g}_k(x) := \frac{\varepsilon_k^2\bar{u}_{2,k}(\mu_{2,k} - \mu_{1,k})}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} &= -\frac{\lambda^4 a_k\bar{u}_{2,k}}{2(a^*)^2\varepsilon_k^2}\int_{\mathbb{R}^2}\frac{\bar{u}_{2,k}^4 - \bar{u}_{1,k}^4}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}}dx \\ &= -\frac{\lambda^4 a_k\bar{u}_{2,k}}{2(a^*)^2\varepsilon_k^2}\int_{\mathbb{R}^2}\bar{\xi}_k\big(\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k} + \bar{u}_{1,k}\big)dx,\end{aligned} \tag{2.26}$$

due to the relation (2.15).

Motivated by [3], we first claim that for any $x_0 \in \mathbb{R}^2$, there exists a small constant $\delta > 0$ such that

$$\int_{\partial B_\delta(x_0)}\Big[\varepsilon_k^2|\nabla\bar{\xi}_k|^2 + \frac{\lambda^2}{2}|\bar{\xi}_k|^2 + \varepsilon_k^2 h(x)|\bar{\xi}_k|^2\Big]dS = O(\varepsilon_k^2) \quad \text{as } k \to \infty. \tag{2.27}$$

To prove the above claim, multiplying (2.24) by $\bar{\xi}_k$ and integrating over $\mathbb{R}^2$, we obtain that

$$\begin{aligned}
&\varepsilon_k^2\int_{\mathbb{R}^2}|\nabla\bar{\xi}_k|^2-\mu_{i,k}\varepsilon_k^2\int_{\mathbb{R}^2}|\bar{\xi}_k|^2+\varepsilon_k^2\int_{\mathbb{R}^2}h(x)|\bar{\xi}_k|^2\\
&=\frac{\lambda^2a_k}{a^*}\int_{\mathbb{R}^2}\big(\bar{u}_{2,k}^2+\bar{u}_{2,k}\bar{u}_{1,k}+\bar{u}_{1,k}^2\big)|\bar{\xi}_k|^2\\
&\quad-\frac{\lambda^4a_k}{2(a^*)^2\varepsilon_k^2}\int_{\mathbb{R}^2}\bar{u}_{2,k}\bar{\xi}_k\int_{\mathbb{R}^2}\bar{\xi}_k\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\\
&\le\frac{\lambda^2a_k}{a^*}\int_{\mathbb{R}^2}\big(\bar{u}_{2,k}^2+\bar{u}_{2,k}\bar{u}_{1,k}+\bar{u}_{1,k}^2\big)+\frac{\lambda^4a_k}{2(a^*)^2\varepsilon_k^2}\int_{\mathbb{R}^2}\bar{u}_{2,k}\int_{\mathbb{R}^2}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\\
&\le C\varepsilon_k^2\ \text{ as }\ k\to\infty,
\end{aligned}$$

due to the uniform boundedness of $|\bar{\xi}_k|$. This implies that there exists a constant $C_1>0$ such that

$$I:=\varepsilon_k^2\int_{\mathbb{R}^2}|\nabla\bar{\xi}_k|^2+\frac{\lambda^2}{2}\int_{\mathbb{R}^2}|\bar{\xi}_k|^2+\varepsilon_k^2\int_{\mathbb{R}^2}h(x)|\bar{\xi}_k|^2<C_1\varepsilon_k^2\quad\text{as }\ k\to\infty. \tag{2.28}$$

Applying Lemma 4.5 in [3], we then conclude that for any $x_0\in\mathbb{R}^2$, there exist a small constant $\delta>0$ and $C_2>0$ such that

$$\int_{\partial B_\delta(x_0)}\Big[\varepsilon_k^2|\nabla\bar{\xi}_k|^2+\frac{\lambda^2}{2}|\bar{\xi}_k|^2+\varepsilon_k^2h(x)|\bar{\xi}_k|^2\Big]dS\le C_2I\le C_1C_2\varepsilon_k^2\ \text{ as }\ k\to\infty,$$

which therefore implies the claim (2.27).

We next define

$$\xi_k(x)=\bar{\xi}_k\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big),\quad k=1,2,\cdots, \tag{2.29}$$

and

$$\tilde{u}_{i,k}(x):=\frac{\sqrt{a^*}\varepsilon_k}{\lambda}u_{i,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big),\quad\text{where }\ i=1,2,$$

so that $\tilde{u}_{i,k}(x)\to w(x)$ uniformly in $\mathbb{R}^2$ as $k\to\infty$ in view of Proposition 2.2. Under the additional assumption (1.11), we shall carry out the proof of Theorem 2.1 by deriving a contradiction through the following four steps.

Step 1. There exist a subsequence $\{a_k\}$ and some constants b_0, b_1 and b_2 such that $\xi_k(x)\to\xi_0(x)$ in $C_{loc}(\mathbb{R}^2)$ as $k\to\infty$, where

$$\xi_0(x)=b_0\big(w+x\cdot\nabla w\big)+\sum_{i=1}^2b_i\frac{\partial w}{\partial x_i}. \tag{2.30}$$

Note that ξ_k satisfies

$$-\Delta\xi_k+C_k(x)\xi_k=g_k(x)\quad\text{in }\ \mathbb{R}^2, \tag{2.31}$$

where the coefficient $C_k(x)$ satisfies

$$\begin{aligned}
C_k(x):=&-\Big(1-\frac{\varepsilon_k^{2+p}}{a^*}\Big)\big(\tilde{u}_{2,k}^2(x)+\tilde{u}_{2,k}(x)\tilde{u}_{1,k}(x)+\tilde{u}_{1,k}^2(x)\big)\\
&-\frac{\varepsilon_k^2}{\lambda^2}\mu_{1,k}+\frac{\varepsilon_k^2}{\lambda^2}h\big(\frac{\varepsilon_kx}{\lambda}+x_{2,k}\big),
\end{aligned} \tag{2.32}$$

and the nonhomogeneous term $g_k(x)$ satisfies

$$\begin{aligned} g_k(x) := \frac{\tilde{u}_{2,k}}{\lambda^2}\frac{\varepsilon_k^2(\mu_{2,k}-\mu_{1,k})}{\|\tilde{u}_{2,k}-\tilde{u}_{1,k}\|_{L^\infty}} &= -\frac{\tilde{u}_{2,k}}{\lambda^2}\frac{a_k\varepsilon_k^2}{2}\int_{\mathbb{R}^2}\frac{u_{2,k}^4-u_{1,k}^4}{\|\tilde{u}_{2,k}-\tilde{u}_{1,k}\|_{L^\infty}}dx \\ &= -\frac{a_k\tilde{u}_{2,k}}{2(a^*)^2}\int_{\mathbb{R}^2}\xi_k\big(\tilde{u}_{2,k}^2+\tilde{u}_{1,k}^2\big)\big(\tilde{u}_{2,k}+\tilde{u}_{1,k}\big)dx. \end{aligned} \tag{2.33}$$

Here we have used (2.15) and (2.23). Since $\|\xi_k\|_{L^\infty(\mathbb{R}^2)} \le 1$, the standard elliptic regularity then implies that $\|\xi_k\|_{C^{1,\alpha}_{loc}(\mathbb{R}^2)} \le C$ for some $\alpha \in (0,1)$, where the constant $C>0$ is independent of k. Therefore, there exist a subsequence $\{a_k\}$ and a function $\xi_0 = \xi_0(x)$ such that $\xi_k(x) \to \xi_0(x)$ in $C_{loc}(\mathbb{R}^2)$ as $k \to \infty$. Applying Proposition 2.2, direct calculations yield from (2.15) and (2.16) that

$$C_k(x) \to 1 - 3w^2(x) \ \text{ uniformly on } \mathbb{R}^2 \ \text{ as } \ k \to \infty,$$

and

$$g_k(x) \to -\frac{2w(x)}{a^*}\int_{\mathbb{R}^2} w^3\xi_0 \ \text{ uniformly on } \mathbb{R}^2 \ \text{ as } \ k \to \infty.$$

This implies that ξ_0 solves

$$-\Delta\xi_0 + (1-3w^2)\xi_0 = \Big(-\frac{2}{a^*}\int_{\mathbb{R}^2} w^3\xi_0\Big)w \ \text{ in } \ \mathbb{R}^2. \tag{2.34}$$

Applying (2.13), we then conclude from (2.34) that (2.30) holds for some constants b_0, b_1 and b_2.

Step 2. The constants b_0, b_1 and b_2 in (2.30) satisfy the following Pohozaev-type system

$$b_0(p-1)\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}w^2 + b_1\int_{\mathbb{R}^2}\frac{\partial^2 h(x)}{\partial x_j\partial x_1}w^2 + b_2\int_{\mathbb{R}^2}\frac{\partial^2 h(x)}{\partial x_j\partial x_2}w^2 = 0, \quad j=1,\,2; \tag{2.35}$$

$$pb_0\int_{\mathbb{R}^2} h(x)w^2 + b_1\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_1}w^2 + b_2\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_2}w^2 = 0. \tag{2.36}$$

We first derive (2.35). Multiplying (2.23) by $\frac{\partial \bar{u}_{i,k}}{\partial x_j}$, where $i,j=1,2$, and integrating over $B_\delta(x_{2,k})$, where $\delta>0$ is small and given by (2.27), we calculate that

$$\begin{aligned} &-\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\Delta\bar{u}_{i,k} + \varepsilon_k^2\int_{B_\delta(x_{2,k})} h(x)\frac{\partial\bar{u}_{i,k}}{\partial x_j}\bar{u}_{i,k} \\ &= \mu_{i,k}\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\bar{u}_{i,k} + \frac{\lambda^2 a_k}{a^*}\int_{B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\bar{u}_{i,k}^3 \\ &= \frac{1}{2}\mu_{i,k}\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^2\nu_j dS + \frac{\lambda^2 a_k}{4a^*}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^4\nu_j dS, \end{aligned} \tag{2.37}$$

where $\nu = (\nu_1,\nu_2)$ denotes the outward unit normal of $\partial B_\delta(x_{2,k})$. Note that

$$\begin{aligned} &-\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\Delta\bar{u}_{i,k} \\ &= -\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\frac{\partial\bar{u}_{i,k}}{\partial\nu}dS + \varepsilon_k^2\int_{B_\delta(x_{2,k})}\nabla\bar{u}_{i,k}\cdot\nabla\frac{\partial\bar{u}_{i,k}}{\partial x_j} \\ &= -\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\frac{\partial\bar{u}_{i,k}}{\partial\nu}dS + \frac{1}{2}\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}|\nabla\bar{u}_{i,k}|^2\nu_j dS, \end{aligned}$$

and

$$\varepsilon_k^2\int_{B_\delta(x_{2,k})}h(x)\frac{\partial\bar{u}_{i,k}}{\partial x_j}\bar{u}_{i,k}=\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}h(x)\bar{u}_{i,k}^2\nu_j dS-\frac{\varepsilon_k^2}{2}\int_{B_\delta(x_{2,k})}\frac{\partial h(x)}{\partial x_j}\bar{u}_{i,k}^2.$$

We then derive from (2.37) that

$$\begin{aligned}
&\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial h(x)}{\partial x_j}\bar{u}_{i,k}^2\\
&=-2\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial x_j}\frac{\partial\bar{u}_{i,k}}{\partial\nu}dS+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}|\nabla\bar{u}_{i,k}|^2\nu_j dS\\
&\quad+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}h(x)\bar{u}_{i,k}^2\nu_j dS-\mu_{i,k}\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^2\nu_j dS\\
&\quad-\frac{\lambda^2 a_k}{2a^*}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^4\nu_j dS.
\end{aligned}\tag{2.38}$$

Following (2.38), we thus have

$$\begin{aligned}
&\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial h(x)}{\partial x_j}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k dx\\
&=-2\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\Big[\frac{\partial\bar{u}_{2,k}}{\partial x_j}\frac{\partial\bar{\xi}_k}{\partial\nu}+\frac{\partial\bar{\xi}_k}{\partial x_j}\frac{\partial\bar{u}_{1,k}}{\partial\nu}\Big]dS\\
&\quad+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\nabla\bar{\xi}_k\cdot\nabla\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\nu_j dS\\
&\quad+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS-\mu_{1,k}\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS\\
&\quad-\frac{\lambda^2 a_k}{2a^*}\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS\\
&\quad-\frac{\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{2,k}^2\nu_j dS.
\end{aligned}\tag{2.39}$$

We now estimate the right hand side of (2.39) as follows. Applying (2.27), if $\delta>0$ is small, we then deduce that

$$\begin{aligned}
&\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\Big|\frac{\partial\bar{u}_{2,k}}{\partial x_j}\frac{\partial\bar{\xi}_k}{\partial\nu}\Big|dS\\
&\le\varepsilon_k\Big(\int_{\partial B_\delta(x_{2,k})}\Big|\frac{\partial\bar{u}_{2,k}}{\partial x_j}\Big|^2dS\Big)^{\frac{1}{2}}\Big(\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\Big|\frac{\partial\bar{\xi}_k}{\partial\nu}\Big|^2dS\Big)^{\frac{1}{2}}\le C\varepsilon_k^2e^{-\frac{C\delta}{\varepsilon_k}}\ \text{ as }\ k\to\infty,
\end{aligned}\tag{2.40}$$

due to the fact that $\nabla\bar{u}_{2,k}$ satisfies the exponential decay (2.20), where $C>0$ is independent of k. Similarly, we have

$$\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\Big|\frac{\partial\bar{\xi}_k}{\partial x_j}\frac{\partial\bar{u}_{1,k}}{\partial\nu}\Big|dS\le C\varepsilon_k^2e^{-\frac{C\delta}{\varepsilon_k}}\ \text{ as }\ k\to\infty,$$

and

$$\varepsilon_k^2\Big|\int_{\partial B_\delta(x_{2,k})}\nabla\bar{\xi}_k\cdot\nabla\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\nu_j dS\Big|\le C\varepsilon_k^2e^{-\frac{C\delta}{\varepsilon_k}}\ \text{ as }\ k\to\infty,$$

On the other hand, since both $|\bar{\xi}_k|$ and $|\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2|$ are bounded uniformly in k, we also get that

$$\begin{aligned}
&\Big|\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})} h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS-\mu_{1,k}\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS\\
&-\frac{\lambda^2 a_k}{2a^*}\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\nu_j dS\\
&-\frac{\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{2,k}^2\nu_j dS\Big|\\
=\,&o(e^{-\frac{C\delta}{\varepsilon_k}})\ \text{ as }\ k\to\infty,
\end{aligned}\tag{2.41}$$

where (2.49) is also used. Because $h(x)$ is homogeneous of degree p, it then follows from (2.39) that for small $\delta>0$,

$$\begin{aligned}
o(e^{-\frac{C\delta}{\varepsilon_k}}) &= \varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{\partial h(x)}{\partial x_j}\big[\bar{u}_{2,k}(x)+\bar{u}_{1,k}(x)\big]\bar{\xi}_k(x)dx\\
&=\frac{\varepsilon_k^3}{\lambda}\int_{B_{\frac{\lambda\delta}{\varepsilon_k}}(0)}\frac{\partial}{\partial y_j}h\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)\bar{\xi}_k\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)\\
&\qquad\cdot\Big[\bar{u}_{2,k}\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)+\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)\Big]dy\\
&=\frac{\varepsilon_k^{p+3}}{\lambda^{p+1}}\int_{B_{\frac{\lambda\delta}{\varepsilon_k}}(0)}\frac{\partial}{\partial y_j}h\big(y+\frac{\lambda x_{2,k}}{\varepsilon_k}\big)\bar{\xi}_k\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)\\
&\qquad\cdot\Big[\bar{u}_{2,k}\big(\frac{\varepsilon_k}{\lambda}y+x_{2,k}\big)+\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}y+x_{1,k}\big)\Big]dy+\varepsilon_k^{p+3}O\big(\frac{x_{2,k}-x_{1,k}}{\varepsilon_k}\big)
\end{aligned}\tag{2.42}$$

as $k\to\infty$. We thus derive from (2.6), (2.30) and (2.42) that

$$\begin{aligned}
0=2\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}w\,\xi_0 &= 2\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}w\Big[b_0\big(w+x\cdot\nabla w\big)+\sum_{i=1}^2 b_i\frac{\partial w}{\partial x_i}\Big]\\
&=b_0\Big[2\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}w^2+\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}\big(x\cdot\nabla w^2\big)\Big]\\
&\quad-\sum_{i=1}^2 b_i\int_{\mathbb{R}^2}\frac{\partial^2 h(x)}{\partial x_j\partial x_i}w^2,\quad j=1,\,2.
\end{aligned}$$

Since $x\cdot\nabla h(x)=ph(x)$, one can check that

$$\begin{aligned}
\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}\big(x\cdot\nabla w^2\big) &= \int_{\mathbb{R}^2}\frac{\partial w^2}{\partial x_j}\big(x\cdot\nabla h(x)\big)+\int_{\mathbb{R}^2}h(x)\frac{\partial w^2}{\partial x_j}\\
&=-(p+1)\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_j}w^2,\quad j=1,\,2,
\end{aligned}$$

and therefore (2.35) follows from above two identities.

We next derive (2.36). Using the integration by parts, we note that

$$\begin{aligned}
&-\varepsilon_k^2\int_{B_\delta(x_{2,k})}\big[(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\big]\Delta\bar{u}_{i,k}\\
&=-\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial\nu}(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}+\varepsilon_k^2\int_{B_\delta(x_{2,k})}\nabla\bar{u}_{i,k}\nabla\big[(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\big] \qquad (2.43)\\
&=-\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial\nu}(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}+\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\big[(x-x_{2,k})\cdot\nu\big]|\nabla\bar{u}_{i,k}|^2.
\end{aligned}$$

Multiplying (2.23) by $(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}$, where $i=1,2$, and integrating over $B_\delta(x_{2,k})$, where $\delta>0$ is small as before, we deduce that for $i=1,2$,

$$\begin{aligned}
&-\varepsilon_k^2\int_{B_\delta(x_{2,k})}\big[(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\big]\Delta\bar{u}_{i,k}\\
&=\varepsilon_k^2\int_{B_\delta(x_{2,k})}\big[\mu_{i,k}-h(x)\big]\bar{u}_{i,k}\big[(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\big]\\
&\quad+\frac{\lambda^2a_k}{a^*}\int_{B_\delta(x_{2,k})}\bar{u}_{i,k}^3\big[(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\big]\\
&=-\frac{\varepsilon_k^2}{2}\int_{B_\delta(x_{2,k})}\bar{u}_{i,k}^2\Big\{2\big[\mu_{i,k}-h(x)\big]-(x-x_{2,k})\cdot\nabla h(x)\Big\} \qquad (2.44)\\
&\quad+\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^2\big[\mu_{i,k}-h(x)\big](x-x_{2,k})\nu dS\\
&\quad-\frac{\lambda^2a_k}{2a^*}\int_{B_\delta(x_{2,k})}\bar{u}_{i,k}^4+\frac{\lambda^2a_k}{4a^*}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^4(x-x_{2,k})\nu dS\\
&=-\mu_{i,k}\varepsilon_k^2\int_{\mathbb{R}^2}\bar{u}_{i,k}^2+\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2}h(x)\bar{u}_{i,k}^2-\frac{\lambda^2a_k}{2a^*}\int_{\mathbb{R}^2}\bar{u}_{i,k}^4+I_i,
\end{aligned}$$

where the lower order term I_i satisfies

$$\begin{aligned}
I_i&=\mu_{i,k}\varepsilon_k^2\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\bar{u}_{i,k}^2-\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}h(x)\bar{u}_{i,k}^2\\
&\quad+\frac{\lambda^2a_k}{2a^*}\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\bar{u}_{i,k}^4-\frac{1}{2}|x_{2,k}|\varepsilon_k^2\int_{B_\delta(x_{2,k})}\bar{u}_{i,k}^2\frac{x_{2,k}}{|x_{2,k}|}\cdot\nabla h(x)\\
&\quad+\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^2\big[\mu_{i,k}-h(x)\big](x-x_{2,k})\nu dS\\
&\quad+\frac{\lambda^2a_k}{4a^*}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{i,k}^4(x-x_{2,k})\nu dS,\ \ i=1,2.
\end{aligned} \qquad (2.45)$$

Since it follows from (2.15) that

$$-\mu_{i,k}\varepsilon_k^2\int_{\mathbb{R}^2}\bar{u}_{i,k}^2-\frac{\lambda^2a_k}{2a^*}\int_{\mathbb{R}^2}\bar{u}_{i,k}^4=-\frac{a^*\varepsilon_k^4}{\lambda^2}\Big[\mu_{i,k}+\frac{a_k}{2}\int_{\mathbb{R}^2}u_{i,k}^4\Big]=-\frac{a^*\varepsilon_k^4}{\lambda^2}e(a_k),$$

we reduce from (2.43)–(2.45) that

$$\begin{aligned}
\tfrac{a^*\varepsilon_k^4}{\lambda^2}e(a_k)-\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2}h(x)\bar{u}_{i,k}^2&=I_i+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\frac{\partial\bar{u}_{i,k}}{\partial\nu}(x-x_{2,k})\cdot\nabla\bar{u}_{i,k}\\
&\quad-\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\big[(x-x_{2,k})\cdot\nu\big]|\nabla\bar{u}_{i,k}|^2,\ \ i=1,2,
\end{aligned}$$

which implies that

$$-\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2}h(x)\big[\bar{u}_{2,k}+\bar{u}_{1,k}\big]\bar{\xi}_k=T_k. \tag{2.46}$$

Here the term T_k satisfies that for small $\delta>0$,

$$\begin{aligned}T_k=&\frac{I_2-I_1}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}-\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\big[(x-x_{2,k})\cdot\nu\big]\big[\nabla\bar{u}_{2,k}+\nabla\bar{u}_{1,k}\big]\bar{\xi}_k\\&+\varepsilon_k^2\int_{\partial B_\delta(x_{2,k})}\Big\{\big[(x-x_{2,k})\cdot\nabla\bar{u}_{2,k}\big]\big(\nu\cdot\nabla\bar{\xi}_k\big)+\big(\nu\cdot\nabla\bar{u}_{1,k}\big)\big[(x-x_{2,k})\cdot\nabla\bar{\xi}_k\big]\Big\}\\=&\frac{I_2-I_1}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}+o(e^{-\frac{C\delta}{\varepsilon_k}})\ \text{ as }\ k\to\infty,\end{aligned} \tag{2.47}$$

due to (2.27), where the second equality follows by applying the argument of estimating (2.40).

Using the arguments of estimating (2.40) and (2.41), along with the exponential decay of $\bar{u}_{i,k}$, we also derive that for small $\delta>0$,

$$\begin{aligned}&\frac{I_2-I_1}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\\=&\mu_{2,k}\varepsilon_k^2\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k-\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\\&+\frac{\lambda^2a_k}{2a^*}\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\\&+\frac{\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\bar{u}_{1,k}^2-\frac{1}{2}|x_{2,k}|\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{x_{2,k}}{|x_{2,k}|}\cdot\nabla h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\\&+\frac{\lambda^2a_k}{4a^*}\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k(x-x_{2,k})\nu dS\\&-\frac{\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_kh(x)(x-x_{2,k})\nu dS\\&+\frac{\mu_{2,k}\varepsilon_k^2}{2}\int_{\partial B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k(x-x_{2,k})\nu dS\\&+\frac{\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2}{2\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{1,k}^2(x-x_{2,k})\nu dS\\=&\frac{\big(\mu_{2,k}-\mu_{1,k}\big)\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\Big[\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\bar{u}_{1,k}^2+\frac{1}{2}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{1,k}^2(x-x_{2,k})\nu dS\Big]\\&-\frac{1}{2}|x_{2,k}|\varepsilon_k^2\int_{B_\delta(x_{2,k})}\frac{x_{2,k}}{|x_{2,k}|}\cdot\nabla h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k+o(e^{-\frac{C\delta}{\varepsilon_k}})\ \text{ as }\ k\to\infty.\end{aligned} \tag{2.48}$$

Since (2.26) gives that

$$\frac{\big|\mu_{2,k}-\mu_{1,k}\big|\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\le\frac{\lambda^4a_k}{2(a^*)^2\varepsilon_k^2}\int_{\mathbb{R}^2}\big(\bar{u}_{2,k}^2+\bar{u}_{1,k}^2\big)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)|\bar{\xi}_k|\le M, \tag{2.49}$$

where the constants $M > 0$ is independent of k, we derive that

$$\frac{(\mu_{2,k}-\mu_{1,k})\varepsilon_k^2}{\|\bar{u}_{2,k}-\bar{u}_{1,k}\|_{L^\infty}}\Big[\int_{\mathbb{R}^2\setminus B_\delta(x_{2,k})}\bar{u}_{1,k}^2+\frac{1}{2}\int_{\partial B_\delta(x_{2,k})}\bar{u}_{1,k}^2(x-x_{2,k})\nu dS\Big]=O(e^{-\frac{C\delta}{\varepsilon_k}}) \quad (2.50)$$

as $k\to\infty$, where the constant $C>0$ is independent of k. Moreover,

$$\begin{aligned}
&\frac{1}{2}|x_{2,k}|\varepsilon_k^2\Big|\int_{B_\delta(x_{2,k})}\frac{x_{2,k}}{|x_{2,k}|}\cdot\nabla h(x)\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)\bar{\xi}_k\Big|\\
&\le\frac{C}{2}\varepsilon_k^3\Big|\frac{x_{2,k}}{\varepsilon_k}\Big|\int_{B_\delta(x_{2,k})}\big(\bar{u}_{2,k}+\bar{u}_{1,k}\big)|\nabla h(x)||\bar{\xi}_k| \qquad (2.51)\\
&=\varepsilon_k^{4+p}O(\Big|\frac{x_{2,k}}{\varepsilon_k}\Big|)=o(\varepsilon_k^{4+p})\ \text{ as }\ k\to\infty,
\end{aligned}$$

where Proposition 2.2 is used for the last equality. Therefore, we deduce from (2.47)–(2.51) that

$$T_k=o(\varepsilon_k^{4+p})\ \text{ as }\ k\to\infty.$$

Further, we obtain from (2.46) that

$$\begin{aligned}
o(\varepsilon_k^{4+p})&=-\frac{2+p}{2}\varepsilon_k^2\int_{\mathbb{R}^2}h(x)\Big[\bar{u}_{2,k}+\bar{u}_{1,k}\Big]\bar{\xi}_k\\
&=-\frac{2+p}{2\lambda^2}\varepsilon_k^4\int_{\mathbb{R}^2}h\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big)\Big[\bar{u}_{2,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big)+\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{1,k}\big)\Big]\xi_k(x)dx\\
&\quad-\frac{2+p}{2\lambda^2}\varepsilon_k^4\int_{\mathbb{R}^2}h\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big)\Big[\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big)-\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{1,k}\big)\Big]\xi_k(x)dx\\
&=-\frac{2+p}{2\lambda^{2+p}}\varepsilon_k^{4+p}\int_{\mathbb{R}^2}h\big(x+\frac{\lambda x_{2,k}}{\varepsilon_k}\big)\Big[\bar{u}_{2,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{2,k}\big)+\bar{u}_{1,k}\big(\frac{\varepsilon_k}{\lambda}x+x_{1,k}\big)\Big]\xi_k(x)dx\\
&\quad+O(\varepsilon_k^{4+p}|x_{2,k}-x_{1,k}|)\ \text{ as }\ k\to\infty.
\end{aligned}$$

By Proposition 2.2 and Step 1, we thus obtain from above that

$$\begin{aligned}
0=2\int_{\mathbb{R}^2}h(x)w\xi_0&=2b_0\int_{\mathbb{R}^2}h(x)w\big(w+x\cdot\nabla w\big)+\sum_{i=1}^2 b_i\int_{\mathbb{R}^2}h(x)\frac{\partial w^2}{\partial x_i}\\
&=2b_0\Big[\int_{\mathbb{R}^2}h(x)w^2+\frac{1}{2}\int_{\mathbb{R}^2}h(x)x\cdot\nabla w^2\Big]-\sum_{i=1}^2 b_i\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_i}w^2\\
&=2b_0\Big\{\int_{\mathbb{R}^2}h(x)w^2-\frac{1}{2}\int_{\mathbb{R}^2}w^2\big[2h(x)+x\cdot\nabla h(x)\big]\Big\}\\
&\quad-\sum_{i=1}^2 b_i\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_i}w^2\\
&=-pb_0\int_{\mathbb{R}^2}h(x)w^2-\sum_{i=1}^2 b_i\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_i}w^2,
\end{aligned}$$

which therefore implies that (2.36) holds.

Step 3. $b_0=b_1=b_2=0$, *i.e.*, $\xi_0\equiv 0$.

One can solve from (2.36) that

$$b_0 = -\frac{1}{p\int_{\mathbb{R}^2} h(x)w^2}\Big[b_1\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_1}w^2 + b_2\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_2}w^2\Big]. \tag{2.52}$$

It then follows from (2.35) and (2.52) that

$$\begin{aligned} &b_1\Big[\int_{\mathbb{R}^2}\frac{\partial^2 h(x)}{\partial x_j\partial x_1}w^2 - \frac{p-1}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_1}w^2\Big] \\ &+b_2\Big[\int_{\mathbb{R}^2}\frac{\partial^2 h(x)}{\partial x_j\partial x_2}w^2 - \frac{p-1}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2}\frac{\partial h(x)}{\partial x_2}w^2\Big] = 0, \quad j=1,\,2. \end{aligned} \tag{2.53}$$

By setting the vector function

$$H(y) = \Big(\int_{\mathbb{R}^2}\Big[\frac{\partial h(x+y)}{\partial y_j} - \frac{p-1}{p\int_{\mathbb{R}^2} h(x)w^2}h(x+y)\Big]w^2(x)dx\Big)_{j=1,2},$$

we have

$$Jac\,H(y) = \Big(\int_{\mathbb{R}^2}\Big[\frac{\partial^2 h(x+y)}{\partial y_j\partial y_i} - \frac{p-1}{p\int_{\mathbb{R}^2} h(x)w^2}\frac{\partial h(x+y)}{\partial y_i}\Big]w^2(x)dx\Big)_{i,j=1,2}.$$

Therefore, if $h(x)$ satisfies (1.11), *i.e.*, if

$$\det\, Jac\,H(0) \neq 0,$$

we then conclude that (2.53) admits the trivial solution $b_1 = b_2 = 0$, which further implies that $b_0 = 0$ in view of (2.52), *i.e.*, $\xi_0 \equiv 0$.

Step 4. $\xi_0 \equiv 0$ cannot occur.

Finally, let y_k be a point satisfying $|\xi_k(y_k)| = \|\xi_k\|_{L^\infty(\mathbb{R}^2)} = 1$. By the same argument as employed in proving Lemma 3.1 in next section, applying the maximum principle to (2.31) yields that $|y_k| \leq C$ uniformly in k. Therefore, we conclude that $\xi_k \to \xi_0 \not\equiv 0$ uniformly on $\mathbb{R}^2$, which however contradicts to the fact that $\xi_0 \equiv 0$ on $\mathbb{R}^2$. This completes the proof of Theorem 2.1. □

3 Refined Spike Profiles

In the following two sections, we shall derive the refined spike profiles of positive minimizers $u_k = u_{a_k}$ for $e(a_k)$ as $a_k \nearrow a^*$. Recall that u_k satisfies the Euler-Lagrange equation (2.14). Under the assumptions of Proposition 2.2, for convenience, we denote

$$\varepsilon_k = (a^* - a_k)^{\frac{1}{2+p}},\ \ \alpha_k := \varepsilon_k^{2+p} \ \text{ and } \ \beta_k := 1 + \frac{\mu_k\varepsilon_k^2}{\lambda^2}, \tag{3.1}$$

where $\mu_k \in \mathbb{R}$ is the Lagrange multiplier of the equation (2.14), so that

$$\alpha_k \to 0 \ \text{ and } \ \beta_k \to 0 \ \text{ as } \ k \to \infty,$$

where (2.17) is used. In order to discuss the refined spike profiles of u_k as $k \to \infty$, the key is thus to obtain the refined estimate of μ_k (equivalently β_k) in terms of ε_k.

We next define

$$w_k(x) := \bar{u}_k(x) - w(x) := \frac{\sqrt{a^*}\varepsilon_k}{\lambda} u_k\Big(\frac{\varepsilon_k}{\lambda}x + x_k\Big) - w(x), \tag{3.2}$$

so that $w_k(x) \to 0$ uniformly in $\mathbb{R}^2$. By applying (2.14), direct calculations then give that $\bar{u}_k$ satisfies

$$-\Delta\bar{u}_k(x) + \frac{\varepsilon_k^2}{\lambda^2}V\big(\frac{\varepsilon_k}{\lambda}x + x_k\big)\bar{u}_k(x) = \frac{\mu_k\varepsilon_k^2}{\lambda^2}\bar{u}_k(x) + \frac{a_k}{a^*}\bar{u}_k^3(x) \text{ in } \mathbb{R}^2.$$

Relating to the operator $\mathcal{L} := -\Delta + (1 - 3w^2)$ in $\mathbb{R}^2$, we also denote the linearized operator

$$\mathcal{L}_k := -\Delta + \big[1 - \big(\bar{u}_k^2 + \bar{u}_k w + w^2\big)\big] \text{ in } \mathbb{R}^2,$$

so that w_k satisfies

$$\begin{aligned}\mathcal{L}_k w_k(x) = -\alpha_k\Big[\frac{1}{a^*}\bar{u}_k^3(x) + \frac{1}{\lambda^{2+p}}g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big)h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k(x)\Big] \\ +\beta_k\bar{u}_k(x) \text{ in } \mathbb{R}^2, \quad \nabla w_k(0) = 0,\end{aligned} \tag{3.3}$$

where the coefficients α_k and β_k are as in (3.1). Define

$$\begin{aligned}\mathcal{L}_k\psi_{1,k}(x) &= -\alpha_k\Big[\frac{1}{\lambda^{2+p}}g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big)h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k(x) \\ &\qquad +\frac{1}{a^*}\bar{u}_k^3(x)\Big] \text{ in } \mathbb{R}^2, \quad \nabla\psi_{1,k}(0) = 0, \\ \mathcal{L}_k\psi_{2,k}(x) &= \beta_k\bar{u}_k(x) \text{ in } \mathbb{R}^2, \quad \nabla\psi_{2,k}(0) = 0,\end{aligned} \tag{3.4}$$

so that the solution $w_k(x)$ of (3.3) satisfies

$$w_k(x) := \psi_{1,k}(x) + \psi_{2,k}(x) \text{ in } \mathbb{R}^2. \tag{3.5}$$

We first employ Proposition 2.2 to address the following estimates of w_k as $k \to \infty$.

Lemma 3.1. *Under the assumptions of Proposition 2.2, we have*

1. *$\psi_{1,k}(x)$ satisfies*

$$\psi_{1,k}(x) = \alpha_k\psi_1(x) + o(\alpha_k) \text{ as } k \to \infty, \tag{3.6}$$

where $\psi_1(x)$ solves uniquely

$$\nabla\psi_1(0) = 0, \quad \mathcal{L}\psi_1(x) = -\frac{1}{a^*}w^3(x) - \frac{g(0)}{\lambda^{2+p}}h(x)w(x) \text{ in } \mathbb{R}^2. \tag{3.7}$$

2. *$\psi_{2,k}(x)$ satisfies*

$$\psi_{2,k}(x) = \beta_k\psi_2(x) + o(\beta_k) \text{ as } k \to \infty, \tag{3.8}$$

where $\psi_2(x)$ solves uniquely

$$\nabla\psi_2(0) = 0, \quad \mathcal{L}\psi_2(x) = w(x) \text{ in } \mathbb{R}^2, \tag{3.9}$$

i.e., $\psi_2(x)$ satisfies

$$\psi_2 = -\frac{1}{2}\big(w + x\cdot\nabla w\big). \tag{3.10}$$

3. w_k satisfies

$$w_k(x) := \alpha_k\psi_1(x) + \beta_k\psi_2(x) + o(\alpha_k + \beta_k) \ \ as \ \ k \to \infty. \tag{3.11}$$

Proof. 1. We first derive $|\psi_{1,k}| \le C\alpha_k$ in $\mathbb{R}^2$ by contradiction. On the contrary, we assume that

$$\lim_{k\to\infty} \frac{\|\psi_{1,k}\|_{L^\infty}}{\alpha_k} = \infty. \tag{3.12}$$

Set $\bar\psi_{1,k} = \frac{\psi_{1,k}}{\|\psi_{1,k}\|_{L^\infty}}$ so that $\|\bar\psi_{1,k}\|_{L^\infty} = 1$. Following (3.4), $\bar\psi_{1,k}$ then satisfies

$$\begin{aligned} &-\Delta\bar\psi_{1,k} + \big[1 - \big(\bar u_k^2 + \bar u_k w + w^2\big)\big]\bar\psi_{1,k} \\ &= -\frac{\alpha_k}{\|\psi_{1,k}\|_\infty}\Big[\frac{1}{\lambda^{2+p}} g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar u_k(x) + \frac{1}{a^*}\bar u_k^3(x)\Big] \ \text{ in } \ \mathbb{R}^2. \end{aligned} \tag{3.13}$$

Let y_k be the global maximum point of $\bar\psi_{1,k}$ so that $\bar\psi_{1,k}(y_k) = \max_{y\in\mathbb{R}^2} \frac{\psi_{1,k}(y)}{\|\psi_{1,k}\|_{L^\infty}} = 1$. Since both $\bar u_k$ and w decay exponentially in view of (2.7), using the maximum principle we derive from (3.13) that $|y_k| \le C$ uniformly in k.

On the other hand, applying the usual elliptic regularity theory, there exists a subsequence, still denoted by $\{\bar\psi_{1,k}\}$, of $\{\bar\psi_{1,k}\}$ such that $\bar\psi_{1,k} \to \bar\psi_1$ weakly in $H^1(\mathbb{R}^2)$ and strongly in $L^q_{loc}(\mathbb{R}^2)$ for all $q \in [2,\infty)$. Here $\bar\psi_1$ satisfies

$$\nabla\bar\psi_1(0) = 0, \quad \mathcal{L}\bar\psi_1(x) = 0 \ \text{ in } \ \mathbb{R}^2,$$

which implies that $\bar\psi_1 = \sum_{i=1}^2 c_i \frac{\partial w}{\partial y_i}$. Since $\nabla\bar\psi_1(0) = 0$, we obtain that $c_1 = c_2 = 0$. Thus, we have $\bar\psi_1(y) \equiv 0$ in $\mathbb{R}^2$, which however contradicts to the fact that $1 = \bar\psi_{1,k}(y_k) \to \bar\psi_1(y_0)$ for some $y_0 \in \mathbb{R}^2$. Therefore, we have $|\psi_{1,k}| \le C\alpha_k$ in $\mathbb{R}^2$.

We next set $\phi_{1,k}(x) = \psi_{1,k}(x) - \alpha_k\psi_1(x)$, where $\psi_1(x)$ is a solution of (3.7). Then either $\phi_{1,k}(x) = O(\alpha_k)$ or $\phi_{1,k}(x) = o(\alpha_k)$ as $k \to \infty$, and $\phi_{1,k}$ satisfies

$$\nabla\phi_{1,k}(0) = 0, \quad -\Delta\phi_{1,k} + \big[1 - \big(\bar u_k^2 + \bar u_k w + w^2\big)\big]\phi_{1,k} = -\alpha_k f_k(x) \ \text{ in } \ \mathbb{R}^2,$$

where $f_k(x)$ satisfies

$$\begin{aligned} f_k(x) = &\big(2w^2 - \bar u_k^2 - \bar u_k w\big)\psi_1(x) + \frac{1}{a^*}\big(\bar u_k^3(x) - w^3(x)\big) \\ &+\frac{1}{\lambda^{2+p}}\Big[g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar u_k(x) - g(0)h(x)w(x)\Big]. \end{aligned}$$

One can note that $f_k(x) \to 0$ uniformly as $k \to \infty$. Therefore, applying the previous argument yields necessarily that $\phi_{1,k}(x) = o(\alpha_k)$ as $k \to \infty$, and the proof of (3.6) is then complete. Also, the property (2.13) gives the uniqueness of solutions for (3.7).

2. Since the proof of (3.8) is very similar to that of (3.6), we omit the details. Further, the property (2.13) gives the uniqueness of ψ_2. Also, one can check directly that $-(w + x\cdot\nabla w)/2$ is a solution of (3.9), which therefore implies that (3.10) holds.

3. The expansion (3.11) now follows immediately from (3.5), (3.6) and (3.8), and the proof is therefore complete. □

3.1 Case of Non-symmetric Potentials

The main aim of this subsection is to establish Theorem 3.6 on the refined spike profiles of u_k as $k \to \infty$, provided that the potential $V(x) = h(x) \in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and is non-symmetric about the origin, *i.e.*, a special case of the assumption (2.1) with $p > 0$. Following (3.1), from now on we denote for simplicity that

$$o\big([\alpha_k + \beta_k]^2\big) = o\big(\alpha_k^2\big) + o\big(\alpha_k\beta_k\big) + o\big(\beta_k^2\big) \ \text{ as } \ k \to \infty, \tag{3.14}$$

where α_k and β_k are defined in (3.1). We first use Lemma 3.1 to establish the following lemmas.

Lemma 3.2. *Suppose $V(x) = h(x) \in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and is not symmetric about the origin. Then there exists an $x_0 \in \mathbb{R}^2$ such that the subsequence $\{x_k\}$ obtained in Proposition 2.2 satisfies*

$$\frac{\alpha_k x_k}{\varepsilon_k} = O\big(\alpha_k^2 x_0\big) + o\big([\alpha_k + \beta_k]^2\big) \ \ \textit{as} \ \ k \to \infty. \tag{3.15}$$

Proof. Multiplying (3.7) and (3.9) by $\frac{\partial w}{\partial x_1}$ and then integrating over $\mathbb{R}^2$, respectively, we obtain from the property (2.13) that

$$\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \mathcal{L} w_k = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} h(x) w = 0. \tag{3.16}$$

Similarly, we derive from (3.3) and (3.11) that

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \mathcal{L}_k w_k &= \beta_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \bar{u}_k - \alpha_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \Big[\frac{1}{a^*} \bar{u}_k^3 + \frac{\bar{u}_k}{\lambda^{2+p}} h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big) \Big] \\ &= \alpha_k \beta_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_1 + o(\alpha_k \beta_k + \beta_k^2) - I_1, \end{aligned} \tag{3.17}$$

where the identity $\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_2 = 0$ is used, due to the radial symmetry of ψ_2. We obtain from (3.16) that

$$\begin{aligned} I_1 &= \alpha_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \Big[\frac{1}{a^*} \bar{u}_k^3 + \frac{\bar{u}_k}{\lambda^{2+p}} h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big) \Big] \\ &= \alpha_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \Big\{ \frac{1}{a^*} \big(\bar{u}_k^3 - w^3\big) + \frac{1}{\lambda^{2+p}} \Big[h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big) \bar{u}_k - h(x) w \Big] \Big\} \\ &= \frac{\alpha_k}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w_k \big(3w^2 + 3w w_k + w_k^2\big) \\ &\quad + \frac{\alpha_k}{\lambda^{2+p}} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \Big[h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big) \bar{u}_k - h(x) w \Big] \\ &= \frac{3\alpha_k^2}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2 \psi_1 + o(\alpha_k^2 + \alpha_k \beta_k) + I_2, \end{aligned} \tag{3.18}$$

where we have used the identity $\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2 \psi_2 = 0$. Further, applying (3.11) and (3.16)

yields that

$$\begin{aligned}\frac{\lambda^{2+p}}{\alpha_k}I_2 &= \int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big\{h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\big[\bar{u}_k-w\big]+\big[h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)-h(x)\big]w\Big\}\\ &= \int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)w_k+o(\alpha_k+\beta_k)\\ &\quad+\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w+o\Big(\frac{x_k}{\varepsilon_k}\Big)\\ &= \alpha_k\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_1+\beta_k\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_2\\ &\quad+\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w+o\Big(\alpha_k+\beta_k+\frac{x_k}{\varepsilon_k}\Big),\end{aligned}\tag{3.19}$$

where (2.6) is used for the second identity. We thus get that

$$\begin{aligned}I_1 = \alpha_k^2\Big[\frac{3}{a^*}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w^2\psi_1+\frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_1\Big]+\frac{\alpha_k\beta_k}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_2\\ +\frac{\alpha_k}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w+o\Big([\alpha_k+\beta_k]^2+\frac{\alpha_k x_k}{\varepsilon_k}\Big).\end{aligned}\tag{3.20}$$

On the other hand, we obtain from (3.16) that

$$\begin{aligned}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\mathcal{L}_k w_k &= \int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\mathcal{L}w_k+\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\big(\mathcal{L}_k-\mathcal{L}\big)w_k\\ &= -\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w_k^2(3w+w_k)\\ &= -3\alpha_k^2\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w\psi_1^2-6\alpha_k\beta_k\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w\psi_1\psi_2+o(\alpha_k^2+\alpha_k\beta_k).\end{aligned}\tag{3.21}$$

Combining (3.17), (3.21) and (3.20), we now conclude from (3.11) that

$$\begin{aligned}&\frac{\alpha_k}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w\\ &= \alpha_k\beta_k\Big[\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\psi_1+6\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w\psi_1\psi_2-\frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_2\Big]\\ &\quad-\alpha_k^2\Big[\frac{3}{a^*}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w^2\psi_1+\frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_1-3\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w\psi_1^2\Big]\\ &\quad+o(\alpha_k^2+\alpha_k\beta_k).\end{aligned}\tag{3.22}$$

We *claim* that the coefficient I_3 of the term $\alpha_k\beta_k$ in (3.22) is zero, *i.e.*,

$$I_3 := \int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\psi_1+6\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}w\psi_1\psi_2-\frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}h(x)\psi_2.\tag{3.23}$$

If (3.23) holds, we then derive from (3.22) that (3.15) holds for some $x_0\in\mathbb{R}^2$, and the lemma is therefore proved.

To complete the proof of the lemma, it next suffices to prove the claim (3.23). Indeed, using the integration by parts, we derive from (3.10) that

$$\begin{aligned}
A :&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + 6\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w\psi_1\psi_2 \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 - 3\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2\psi_1 - \frac{3}{2}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 (x\cdot\nabla w^2) \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 - 3\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2\psi_1 \\
&\quad + \frac{3}{2}\int_{\mathbb{R}^2} w^2\Big[2\frac{\partial w}{\partial x_1}\psi_1 + x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\psi_1\Big)\Big] \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{3}{2}\int_{\mathbb{R}^2} w^2\Big[\frac{\partial w}{\partial x_1}(x\cdot\nabla\psi_1) + \psi_1 x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big].
\end{aligned}$$

Since $x\cdot\nabla h(x) = ph(x)$, we obtain from (3.10) and (3.16) that

$$\begin{aligned}
B :&= -\frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} h(x)\psi_2 \\
&= \frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} h(x)(w + x\cdot\nabla w) \\
&= \frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} h(x) w\frac{\partial w}{\partial x_1} - \frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} w\Big[2h(x)\frac{\partial w}{\partial x_1} + x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}h(x)\Big)\Big] \\
&= -\frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} w\Big\{\big[x\cdot\nabla h(x)\big]\frac{\partial w}{\partial x_1} + h(x)x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big\} \\
&= -\frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} wh(x)\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big].
\end{aligned}$$

By above calculations, we then get from (3.23) that

$$\begin{aligned}
I_3 = A + B = &\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{3}{2}\int_{\mathbb{R}^2} w^2\frac{\partial w}{\partial x_1}(x\cdot\nabla\psi_1) \\
&+ \frac{1}{2}\int_{\mathbb{R}^2}\Big[3w^2\psi_1 - \frac{wh(x)}{\lambda^{2+p}}\Big]\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big].
\end{aligned} \tag{3.24}$$

Applying the integration by parts, we derive from (3.7) that

$$\begin{aligned}
&\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{1}{2}\int_{\mathbb{R}^2}\Big[3w^2\psi_1 - \frac{wh(x)}{\lambda^{2+p}}\Big]\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{1}{2}\int_{\mathbb{R}^2}\Big[3w^2\psi_1 - \Big(\frac{w^3}{a^*} + \frac{wh(x)}{\lambda^{2+p}}\Big)\Big]\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{1}{2}\int_{\mathbb{R}^2}\big(-\Delta\psi_1 + \psi_1\big)\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] \\
&= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 + \frac{1}{2}\int_{\mathbb{R}^2}\big(-\Delta\psi_1\big)\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] - \frac{1}{2}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\big[2\psi_1 + x\cdot\nabla\psi_1\big] \\
&= \frac{1}{2}\int_{\mathbb{R}^2}\big(-\Delta\psi_1\big)\Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] - \frac{1}{2}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}(x\cdot\nabla\psi_1),
\end{aligned}$$

which then gives that

$$\begin{aligned} -2I_3 = -2(A+B) &= \int_{\mathbb{R}^2} \Delta\psi_1 \Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] + \int_{\mathbb{R}^2} \frac{\partial}{\partial x_1}\Big[w - w^3\Big](x\cdot\nabla\psi_1) \\ &= \int_{\mathbb{R}^2} \Delta\psi_1 \Big[x\cdot\nabla\Big(\frac{\partial w}{\partial x_1}\Big)\Big] + \int_{\mathbb{R}^2} \frac{\partial \Delta w}{\partial x_1}(x\cdot\nabla\psi_1). \end{aligned} \tag{3.25}$$

To further simplify I_3, we next rewrite ψ_1 as $\psi_1(x) = \psi_1(r,\theta)$, where $x = r(\cos\theta, \sin\theta)$ and (r,θ) is the polar coordinate in $\mathbb{R}^2$. We then follow from (3.7) and (3.25) that

$$\begin{aligned} -2I_3 &= \int_0^\infty\int_0^{2\pi} \Big\{\big[r\big(\psi_1\big)_r\big]_r + \frac{1}{r}\big(\psi_1\big)_{\theta\theta}\Big\} r\frac{\partial}{\partial r}\big(w'\cos\theta\big)d\theta dr \\ &\quad + \int_0^\infty\int_0^{2\pi} \frac{\partial}{\partial r}\Big(w'' + \frac{w'}{r}\Big)\cos\theta\, r^2\big(\psi_1\big)_r d\theta dr \\ &= -\int_0^\infty\int_0^{2\pi} r\big(\psi_1\big)_r (rw'')'\cos\theta d\theta dr \\ &\quad + \int_0^\infty\int_0^{2\pi} r\big(\psi_1\big)_r\Big[r\frac{\partial}{\partial r}\Big(w'' + \frac{w'}{r}\Big)\Big]\cos\theta d\theta dr \\ &\quad + \int_0^\infty\int_0^{2\pi} \big(\psi_1\big)_{\theta\theta} w''\cos\theta d\theta dr \\ &= -\int_0^\infty\int_0^{2\pi} r\big(\psi_1\big)_r\Big\{(rw'')' - \Big[r\frac{\partial}{\partial r}\Big(w'' + \frac{w'}{r}\Big)\Big]\Big\}\cos\theta d\theta dr \\ &\quad - \int_0^\infty\int_0^{2\pi} \psi_1 w''\cos\theta d\theta dr \\ &= -\int_0^\infty\int_0^{2\pi} \big(\psi_1\big)_r w'\cos\theta d\theta dr - \int_0^\infty\int_0^{2\pi} \psi_1 w''\cos\theta d\theta dr = 0, \end{aligned} \tag{3.26}$$

i.e., $I_3 = 0$, which therefore implies the claim (3.23). □

Remark 3.1. *Whether the point $x_0 \in \mathbb{R}^2$ in Lemma 3.2 is the origin or not is determined completely by the fact that whether the coefficient I_4 of the term α_k^2 in (3.22) is zero or not, where I_4 satisfies*

$$I_4 := \frac{3}{a^*}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2\psi_1 + \frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} h(x)\psi_1 - 3\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w\psi_1^2.$$

One can check from above that I_4 can be further simplified as

$$I_4 := \int_{\mathbb{R}^2} \big(-\Delta\psi_1\big)\frac{\partial \psi_1}{\partial x_1} - \frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2} w\psi_1 \frac{\partial h(x)}{\partial x_1}. \tag{3.27}$$

Under the non-symmetry assumption of $h(x)$, in general it however seems difficult to derive from (3.27) that whether $I_4 = 0$ or not.

Lemma 3.3. *Suppose $V(x) = h(x) \in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and is not symmetric about the origin. Then we have*

$$w_k := \alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 + \alpha_k\beta_k\psi_5 + o([\alpha_k + \beta_k]^2) \quad as \quad k\to\infty, \tag{3.28}$$

where $\psi_1(x)$ and $\psi_2(x)$ are given in Lemma 3.1, and $\psi_i(x)$, $i=3,4,5$, solves uniquely

$$\nabla\psi_i(0)=0 \ \text{ and } \ \mathcal{L}\psi_i(x)=f_i(x) \ \text{ in } \ \mathbb{R}^2, \ \ i=3,4,5, \tag{3.29}$$

and $f_i(x)$ satisfies for some $y_0\in\mathbb{R}^2$,

$$f_i(x)=\begin{cases} 3w\psi_1^2-\Big(\dfrac{3w^2}{a^*}+\dfrac{h(x)}{\lambda^{2+p}}\Big)\psi_1-\dfrac{w}{\lambda^{1+p}}\big(y_0\cdot\nabla h(x)\big), & \text{if } \ i=3;\\ 3w\psi_2^2+\psi_2, & \text{if } \ i=4;\\ 6w\psi_1\psi_2+\psi_1-\Big(\dfrac{3w^2}{a^*}+\dfrac{h(x)}{\lambda^{2+p}}\Big)\psi_2, & \text{if } \ i=5. \end{cases} \tag{3.30}$$

Moreover, ψ_4 is radially symmetric.

Proof. Following Lemma 3.1(3), set

$$v_k=w_k-\alpha_k\psi_1-\beta_k\psi_2,$$

so that

$$\begin{aligned} \mathcal{L}_k w_k &= \mathcal{L}_k(v_k+\alpha_k\psi_1+\beta_k\psi_2)\\ &= \mathcal{L}_k v_k+\alpha_k(\mathcal{L}_k-\mathcal{L})\psi_1+\beta_k(\mathcal{L}_k-\mathcal{L})\psi_2+\alpha_k\mathcal{L}\psi_1+\beta_k\mathcal{L}\psi_2\\ &= \mathcal{L}_k v_k-w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)-\alpha_k\Big[\frac{w^3}{a^*}+\frac{h(x)w}{\lambda^{2+p}}\Big]+\beta_k w. \end{aligned} \tag{3.31}$$

Applying (3.3), we then have

$$\begin{aligned} \mathcal{L}_k v_k &= \mathcal{L}_k w_k+w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)+\alpha_k\Big[\frac{w^3}{a^*}+\frac{h(x)w}{\lambda^{2+p}}\Big]-\beta_k w\\ &= w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)+\beta_k(\bar u_k-w)\\ &\quad -\alpha_k\Big\{\frac{1}{a^*}(\bar u_k^3-w^3)+\frac{1}{\lambda^{2+p}}\Big[h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\bar u_k-h(x)w\Big]\Big\}\\ &= w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)+\beta_k w_k-I_3, \end{aligned} \tag{3.32}$$

where I_3 satisfies

$$\begin{aligned} I_3 &= \frac{\alpha_k}{a^*}w_k(3w^2+3ww_k+w_k^2)\\ &\quad +\frac{\alpha_k}{\lambda^{2+p}}\Big\{h(x)(\bar u_k-w)+\Big[h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)-h(x)\Big]\bar u_k\Big\}\\ &= \frac{\alpha_k}{a^*}w_k(3w^2+3ww_k+w_k^2)+\frac{\alpha_k}{\lambda^{2+p}}\Big[h(x)w_k+\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)\bar u_k\Big]+o\Big(\frac{\alpha_k x_k}{\varepsilon_k}\Big)\\ &= \alpha_k w_k\Big(\frac{3w^2}{a^*}+\frac{h(x)}{\lambda^{2+p}}\Big)+\frac{\alpha_k}{a^*}w_k^2(3w+w_k)\\ &\quad +\frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)\bar u_k+o\Big(\frac{\alpha_k x_k}{\varepsilon_k}\Big). \end{aligned}$$

Using Lemma 3.2, there exists $y_0\in\mathbb{R}^2$ such that

$$\frac{x_k}{\alpha_k\varepsilon_k}\to y_0+o\Big(\frac{[\alpha_k+\beta_k]^2}{\alpha_k^2}\Big) \ \text{ as } \ k\to\infty.$$

We thus obtain from above that

$$\begin{aligned}\mathcal{L}_k v_k &= w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)+\beta_k w_k-\alpha_k w_k\Big(\frac{3w^2}{a^*}+\frac{h(x)}{\lambda^{2+p}}\Big)\\ &\quad-\frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)\bar{u}_k-\frac{\alpha_k}{a^*}w_k^2(3w+w_k)+o\Big(\frac{\alpha_k x_k}{\varepsilon_k}\Big)\\ &=\alpha_k^2\Big[3w\psi_1^2-\Big(\frac{3w^2}{a^*}+\frac{h(x)}{\lambda^{2+p}}\Big)\psi_1-\frac{w}{\lambda^{1+p}}\big(y_0\cdot\nabla h(x)\big)\Big]\\ &\quad+\alpha_k\beta_k\Big[6w\psi_1\psi_2+\psi_1-\Big(\frac{3w^2}{a^*}+\frac{h(x)}{\lambda^{2+p}}\Big)\psi_2\Big]\\ &\quad+\beta_k^2(3w\psi_2^2+\psi_2)+o([\alpha_k+\beta_k]^2)\quad\text{in }\ \mathbb{R}^2.\end{aligned}\tag{3.33}$$

Following (3.33), the same argument of proving Lemma 3.1 then gives (3.28). Finally, since $f_4(x)$ is radially symmetric, there exists a radial solution ψ_4. Further, the property (2.13) gives the uniqueness of ψ_4. Therefore, ψ_4 must be radially symmetric, and the proof is complete. □

Lemma 3.4. *Suppose $V(x)=h(x)\in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and is not symmetric about the origin. Then we have*

$$\int_{\mathbb{R}^2} w\psi_1=0,\quad \int_{\mathbb{R}^2} w\psi_2=0,\tag{3.34}$$

and

$$I=\int_{\mathbb{R}^2}\big(2w\psi_4+\psi_2^2\big)=0.\tag{3.35}$$

However, we have

$$II=2\int_{\mathbb{R}^2} w\psi_5+2\int_{\mathbb{R}^2}\psi_1\psi_2=-\frac{2+p}{2}<0.\tag{3.36}$$

Here $\psi_1(x),\cdots,\psi_5(x)$ are given in Lemmas 3.1 and 3.3.

Since the proof of Lemma 3.4 is very involved, we leave it to the appendix. Following above lemmas, we are now ready to derive the comparison relation between β_k and α_k.

Proposition 3.5. *Suppose $V(x)=h(x)$ is homogeneous of degree p and is not symmetric about the origin. Then there exists a constant $C^*\in\mathbb{R}^+\cup\mathbb{R}^-$ such that*

$$\beta_k=C^*\alpha_k\ \ \text{as}\ \ k\to\infty,\tag{3.37}$$

where C^* satisfies

$$C^*=\frac{2}{2+p}\Big(2\int_{\mathbb{R}^2} w\psi_3+\int_{\mathbb{R}^2}\psi_1^2\Big).\tag{3.38}$$

Moreover, w_k satisfies

$$w_k:=\big[\psi_1+C^*\psi_2\big]\alpha_k+\big[\psi_3+(C^*)^2\psi_4+C^*\psi_5\big]\alpha_k^2+o(\alpha_k^2)\ \ \text{as}\ \ k\to\infty,\tag{3.39}$$

Here $\psi_1(x),\cdots,\psi_5(x)$ are given in Lemmas 3.1 and 3.3.

Proof. Note from (3.2) that w_k satisfies

$$\int_{\mathbb{R}^2} w^2 = \int_{\mathbb{R}^2} \bar{u}_k^2 = \int_{\mathbb{R}^2} \big(w + w_k\big)^2, \ i.e. \ \ 2\int_{\mathbb{R}^2} w w_k + \int_{\mathbb{R}^2} w_k^2 = 0. \tag{3.40}$$

Applying (3.40), we then derive from Lemma 3.3 that

$$\begin{aligned}
0 &= 2\int_{\mathbb{R}^2} w w_k + \int_{\mathbb{R}^2} w_k^2 \\
&= 2\int_{\mathbb{R}^2} w(\alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 + \alpha_k\beta_k\psi_5) \\
&\quad + \int_{\mathbb{R}^2} (\alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 + \alpha_k\beta_k\psi_5)^2 + o([\alpha_k + \beta_k]^2) \\
&= \alpha_k\Big(2\int_{\mathbb{R}^2} w\psi_1\Big) + \beta_k\Big(2\int_{\mathbb{R}^2} w\psi_2\Big) + \beta_k^2\Big(2\int_{\mathbb{R}^2} w\psi_4 + \int_{\mathbb{R}^2}\psi_2^2\Big) \\
&\quad + \alpha_k\beta_k\Big(2\int_{\mathbb{R}^2} w\psi_5 + 2\int_{\mathbb{R}^2}\psi_1\psi_2\Big) + \alpha_k^2\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) + o([\alpha_k + \beta_k]^2) \\
&= -\frac{2+p}{2}\alpha_k\beta_k + \alpha_k^2\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) + o([\alpha_k + \beta_k]^2),
\end{aligned} \tag{3.41}$$

where Lemma 3.4 is used in the last equality. It then follows from (3.41) that

$$2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2 \neq 0,$$

and moreover,

$$-\frac{2+p}{2}\beta_k + \alpha_k\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) = 0, \quad i.e., \ \beta_k = C^*\alpha_k,$$

where $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is as in (3.38). Finally, the expansion (3.39) follows directly from (3.37) and Lemma 3.3, and we are done. □

Lemma 3.2 and Proposition 3.5 give that the subsequence $\{x_k\}$ satisfies the estimate

$$\frac{x_k}{\varepsilon_k} = O(\alpha_k x_0) + o(\alpha_k) \ \text{ as } \ k \to \infty \tag{3.42}$$

for some $x_0 \in \mathbb{R}^2$. We also conclude from (3.1) and Proposition 3.5 that the Lagrange multiplier $\mu_k \in \mathbb{R}$ of the Euler-Lagrange equation (2.14) satisfies

$$\mu_k = -\frac{\lambda}{\varepsilon_k^2} + \lambda^2 C^* \varepsilon_k^p + o(\varepsilon_k^p) \ \text{ as } \ k \to \infty, \tag{3.43}$$

where $\lambda > 0$ is defined by (2.2), $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is given by (3.38). Moreover, following above results we finally conclude the following refined spike profiles.

Theorem 3.6. *Suppose $V(x) = h(x) \in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree p and is not symmetric about the origin, and let u_a be a positive minimizer of (1.1) for $a < a^*$. Then for any sequence $\{a_k\}$ with $a_k \nearrow a^*$ as $k \to \infty$, there exist a subsequence, still*

denoted by $\{a_k\}$*, of* $\{a_k\}$ *and* $\{x_k\} \subset \mathbb{R}^2$ *such that the subsequence* $\{u_k\}$ *satisfies for* $\varepsilon_k := (a^* - a_k)^{\frac{1}{2+p}}$,

$$\begin{aligned} u_k(x) = \frac{\lambda}{\|w\|_2}\Big\{\frac{1}{\varepsilon_k} w\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+p}\Big[\psi_1 + C^*\psi_2\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) \\ + \varepsilon_k^{3+2p}\Big[\psi_3 + (C^*)^2\psi_4 + C^*\psi_5\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big)\Big\} + o(\varepsilon_k^{3+2p}) \ \ as \ \ k \to \infty \end{aligned} \tag{3.44}$$

uniformly in $\mathbb{R}^2$*, and* $\{x_k\}$ *satisfies*

$$\frac{x_k}{\varepsilon_k^{3+p}} = O(y_0) \ \ as \ \ k \to \infty \tag{3.45}$$

for some $y_0 \in \mathbb{R}^2$*. Here* $\psi_1(x), \cdots, \psi_5(x)$ *are given in Lemmas 3.1 and 3.3.*

Proof. The refined spike profile (3.44) follows immediately from (3.2) and (3.39). Also, Lemma 3.2 gives directly the estimate (3.45). □

4 Refined Spike Profiles under Symmetric Potentials

The main purpose of this section is to derive Theorem 4.4 on the refined spike profiles of positive minimizers u_k as $k \to \infty$ for the case where $V(x)$ admits the symmetry. Throughout this section, for generality we shall always assume that $V(x)$ satisfies

(*V*). $V(x) = g(x)h(x)$, where $h(x) \in C^1_{loc}(\mathbb{R}^2)$ is homogeneous of degree $p \in \mathbb{R}^+$ and is symmetric about the origin, and $g(x) \in C^m_{loc}(\mathbb{R}^2)$ for some $m \in \mathbb{N} \cup \{+\infty\}$ satisfies $0 < C \le g(x) \le \frac{1}{C}$ in $\mathbb{R}^2$ and $G(x) := g(x) - g(0)$,

$$\mathcal{D}^\alpha G(0) = 0 \ \text{ for all } \ |\alpha| \le m-1, \ \text{ and } \ \mathcal{D}^\alpha G(0) \neq 0 \ \text{ for some } \ |\alpha| = m,$$

where it takes $m = +\infty$ if $g(x) \equiv 1$. Recall that Lemma 3.1 still holds for such types of potentials $V(x)$, and let α_k and β_k be defined in (3.1). Similar to Lemma 3.2, we start with the following estimates.

Lemma 4.1. *Suppose* $V(x) = g(x)h(x)$ *satisfies the assumption* (*V*) *for* $p > 0$ *and* $m \in \mathbb{N} \cup \{+\infty\}$*. Then the subsequence* $\{x_k\}$ *obtained in Proposition 2.2 satisfies the following estimates:*

1. *If* m *is even, then we have*

$$\frac{\lambda\alpha_k x_k}{\varepsilon_k} = o\big([\alpha_k + \beta_k]^2 + \alpha_k\varepsilon_k^m\big) \ \ as \ \ k \to \infty. \tag{4.1}$$

2. *If* m *is odd, then we have*

$$\frac{\lambda\alpha_k x_k}{\varepsilon_k} = O(\alpha_k\varepsilon_k^m x_0) + o\big([\alpha_k + \beta_k]^2 + \alpha_k\varepsilon_k^m\big) \ \ as \ \ k \to \infty, \tag{4.2}$$

where $x_0 \in \mathbb{R}^2$ *satisfies*

$$g(0)\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\big[x_0 \cdot \nabla h(x)\big]w + \frac{1}{\lambda^m}\sum_{|\alpha|=m}\int_{\mathbb{R}^2}\frac{\partial w}{\partial x_1}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w = 0. \tag{4.3}$$

Proof. Recall that $\psi_1(x)$ and $\psi_2(x)$ are given in Lemma 3.1. Under the assumptions of $h(x)$, since ψ_1 is symmetric about the origin, we have

$$\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\psi_1 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w\psi_1^2 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w\psi_1\psi_2 = 0.$$

Using the symmetry of ψ_1 and ψ_2, the same calculations of (3.17)–(3.18) yield that

$$\begin{aligned}
o(\alpha_k^2 + \alpha_k\beta_k) &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\mathcal{L}_k w_k \\
&= o(\alpha_k\beta_k + \beta_k^2) - \alpha_k \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[\frac{1}{a^*}\bar{u}_k^3 + \frac{\bar{u}_k}{\lambda^{2+p}} g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\Big] \\
&= o(\alpha_k\beta_k + \beta_k^2) - \frac{\alpha_k}{a^*}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\big(\bar{u}_k^3 - w^3\big) \\
&\quad - \frac{\alpha_k}{\lambda^{2+p}}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k - g(0)h(x)w\Big] \\
&= o(\alpha_k\beta_k + \beta_k^2) - \frac{\alpha_k}{\lambda^{2+p}}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k - g(0)h(x)w\Big] \\
&= o(\alpha_k\beta_k + \beta_k^2) - I_1.
\end{aligned} \tag{4.4}$$

By the assumption (V), similar to (3.19) we get that

$$\begin{aligned}
\frac{\lambda^{2+p}}{\alpha_k} I_1 &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big\{g(0)h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\big[\bar{u}_k - w\big] + g(0)\big[h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big) - h(x)\big]w\Big\} \\
&\quad + \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big) - g(0)\Big] h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k \\
&= o(\alpha_k + \beta_k) + g(0)\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w \\
&\quad + \Big(\frac{\varepsilon_k}{\lambda}\Big)^m \sum_{|\alpha|=m}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[\frac{1}{\alpha!}\Big(x + \frac{\lambda x_k}{\varepsilon_k}\Big)^\alpha \mathcal{D}^\alpha g(0)\Big] h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k + o(\varepsilon_k^m),
\end{aligned}$$

which then implies that

$$\begin{aligned}
I_1 &= \frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\varepsilon_k}{\lambda}\Big)^m \sum_{|\alpha|=m}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[\frac{1}{\alpha!}\Big(x + \frac{\lambda x_k}{\varepsilon_k}\Big)^\alpha \mathcal{D}^\alpha g(0)\Big] h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k \\
&\quad + \frac{\alpha_k}{\lambda^{2+p}} g(0)\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w + o(\alpha_k^2 + \alpha_k\beta_k + \alpha_k\varepsilon_k^m).
\end{aligned} \tag{4.5}$$

Combining (4.4) and (4.5), we then conclude from the estimate (3.11) that

$$\begin{aligned}
&\frac{\alpha_k}{\lambda^{2+p}} g(0)\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)w \\
&= -\frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\varepsilon_k}{\lambda}\Big)^m \sum_{|\alpha|=m}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x)w + o\Big(\big[\alpha_k + \beta_k\big]^2 + \alpha_k\varepsilon_k^m\Big).
\end{aligned} \tag{4.6}$$

If m is even, one can note that

$$\sum_{|\alpha|=m}\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x)w = 0,$$

and it then follows from (4.6) that (4.1) holds. If m is odd, we then derive from (4.6) that both (4.2) and (4.3) hold. □

Lemma 4.2. *Suppose $V(x) = g(x)h(x)$ satisfies the assumption (V) for $p > 0$ and $m \in \mathbb{N} \cup \{+\infty\}$, and let $\psi_1(x)$ and $\psi_2(x)$ be given in Lemma 3.1. Then w_k satisfies*

$$\begin{aligned} w_k :=& \alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 \\ &+\alpha_k\varepsilon_k^m\phi + \alpha_k\beta_k\psi_5 + o\big([\alpha_k + \beta_k]^2 + \alpha_k\varepsilon_k^m\big) \quad as \quad k \to \infty, \end{aligned} \tag{4.7}$$

where $\psi_i(x)$, $i = 3, 4, 5$, solves uniquely

$$\nabla\psi_i(0) = 0 \ \ and \ \ \mathcal{L}\psi_i(x) = g_i(x) \ \ in \ \ \mathbb{R}^2, \ \ i = 3, 4, 5, \tag{4.8}$$

and $g_i(x)$ satisfies

$$g_i(x) = \begin{cases} 3w\psi_1^2 - \Big(\dfrac{3w^2}{a^*} + \dfrac{g(0)h(x)}{\lambda^{2+p}}\Big)\psi_1, & if \ \ i = 3; \\ 3w\psi_2^2 + \psi_2, & if \ \ i = 4; \\ 6w\psi_1\psi_2 + \psi_1 - \Big(\dfrac{3w^2}{a^*} + \dfrac{g(0)h(x)}{\lambda^{2+p}}\Big)\psi_2, & if \ \ i = 5. \end{cases} \tag{4.9}$$

Here ϕ solves uniquely

$$\begin{aligned} \mathcal{L}\phi(x) = -\frac{1}{\lambda^{2+p}}\Big\{ &\big[x_0 \cdot \nabla h(x)\big]g(0)w \\ &+\frac{1}{\lambda^m}\sum_{|\alpha|=m}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w\Big\} \ \ in \ \ \mathbb{R}^2, \ \ and \ \ \nabla\phi(0) = 0, \end{aligned} \tag{4.10}$$

where $x_0 = 0$ for the case where m is even, and $x_0 \in \mathbb{R}^2$ is given by (4.3) for the case where m is odd.

Proof. Following Lemma 3.1(3), we set

$$v_k = w_k - \alpha_k\psi_1 - \beta_k\psi_2.$$

Similar to (3.32), we then have

$$\begin{aligned} \mathcal{L}_k v_k &= w_k(\alpha_k\psi_1 + \beta_k\psi_2)(3w + w_k) + \beta_k w_k - \frac{\alpha_k}{a^*}(\bar{u}_k^3 - w^3) \\ &\quad -\frac{\alpha_k}{\lambda^{2+p}}\Big[g\big(\frac{\varepsilon_k x}{\lambda} + x_k\big)h\big(x + \frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k - g(0)h(x)w\Big] \\ &= w_k(\alpha_k\psi_1 + \beta_k\psi_2)(3w + w_k) + \beta_k w_k \\ &\quad -\frac{\alpha_k}{a^*}w_k(3w^2 + 3ww_k + w_k^2) - I_2, \end{aligned} \tag{4.11}$$

where I_2 satisfies

$$\begin{aligned}
I_2 &= \frac{\alpha_k}{\lambda^{2+p}}\Big\{\Big[g\big(\frac{\varepsilon_k x}{\lambda}+x_k\big)-g(0)\Big]h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k \\
&\qquad +g(0)\Big[h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)-h(x)\Big]\bar{u}_k+g(0)h(x)\big(\bar{u}_k-w\big)\Big\} \\
&= \frac{\alpha_k}{\lambda^{2+p}}\Big\{\Big(\frac{\varepsilon_k}{\lambda}\Big)^m\sum_{|\alpha|=m}\Big[\frac{1}{\alpha!}\Big(x+\frac{\lambda x_k}{\varepsilon_k}\Big)^\alpha\mathcal{D}^\alpha g(0)\Big]h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k \\
&\qquad +g(0)\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)\bar{u}_k+g(0)h(x)w_k\Big\}+o\Big(\frac{\alpha_k x_k}{\varepsilon_k}+\alpha_k\varepsilon_k^m\Big) \\
&= \alpha_k w_k\frac{g(0)h(x)}{\lambda^{2+p}}+\frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)g(0)\bar{u}_k \\
&\quad +\frac{\alpha_k\varepsilon_k^m}{\lambda^{2+p+m}}\sum_{|\alpha|=m}\Big[\frac{1}{\alpha!}\Big(x+\frac{\lambda x_k}{\varepsilon_k}\Big)^\alpha\mathcal{D}^\alpha g(0)\Big]h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k+o\big(\alpha_k\varepsilon_k^m\big),
\end{aligned}\tag{4.12}$$

where Lemma 4.1 is used. Applying Lemma 4.1 again, we then obtain from (4.11) and (4.12) that

$$\begin{aligned}
\mathcal{L}_k v_k &= w_k(\alpha_k\psi_1+\beta_k\psi_2)(3w+w_k)+\beta_k w_k-\frac{\alpha_k}{a^*}w_k^2(3w+w_k) \\
&\quad -\alpha_k w_k\Big[\frac{3w^2}{a^*}+\frac{g(0)h(x)}{\lambda^{2+p}}\Big]-\frac{\alpha_k}{\lambda^{2+p}}\Big(\frac{\lambda x_k}{\varepsilon_k}\cdot\nabla h(x)\Big)g(0)\bar{u}_k \\
&\quad -\frac{\alpha_k\varepsilon_k^m}{\lambda^{2+p+m}}\sum_{|\alpha|=m}\Big[\frac{1}{\alpha!}\Big(x+\frac{\lambda x_k}{\varepsilon_k}\Big)^\alpha\mathcal{D}^\alpha g(0)\Big]h\big(x+\frac{\lambda x_k}{\varepsilon_k}\big)\bar{u}_k+o\big(\alpha_k\varepsilon_k^m\big) \\
&= \alpha_k^2\Big[3w\psi_1^2-\Big(\frac{3w^2}{a^*}+\frac{g(0)h(x)}{\lambda^{2+p}}\Big)\psi_1\Big] \\
&\quad +\alpha_k\beta_k\Big[6w\psi_1\psi_2+\psi_1-\Big(\frac{3w^2}{a^*}+\frac{g(0)h(x)}{\lambda^{2+p}}\Big)\psi_2\Big] \\
&\quad -\frac{\alpha_k\varepsilon_k^m}{\lambda^{2+p}}\Big\{\big[x_0\cdot\nabla h(x)\big]g(0)w+\frac{1}{\lambda^m}\sum_{|\alpha|=m}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w\Big\} \\
&\quad +\beta_k^2(3w\psi_2^2+\psi_2)+o\big([\alpha_k+\beta_k]^2+\alpha_k\varepsilon_k^m\big)\quad\text{in }\ \mathbb{R}^2,
\end{aligned}\tag{4.13}$$

where $x_0=0$ for the case where m is even, and $x_0\in\mathbb{R}^2$ is given by (4.3) for the case where m is odd. Following (4.13), the same argument of proving Lemma 3.1 then gives (4.7), and the proof is therefore complete. □

Proposition 4.3. *Suppose $V(x)=g(x)h(x)$ satisfies the assumption (V) for $p>0$ and $m\in\mathbb{N}\cup\{+\infty\}$. Let $\psi_1(x),\cdots,\psi_5(x)$ be given in Lemmas 3.1 and 4.2, and ϕ is given by (4.10).*

1. *If $m>2+p$, then*

$$\beta_k=C^*\alpha_k,\tag{4.14}$$

and w_k satisfies

$$w_k:=\big[\psi_1+C^*\psi_2\big]\alpha_k+\big[\psi_3+(C^*)^2\psi_4+C^*\psi_5\big]\alpha_k^2+o(\alpha_k^2)\ \ as\ \ k\to\infty,\tag{4.15}$$

where the constant $C^*\in\mathbb{R}^+\cup\mathbb{R}^-$ is given by (3.38).

2. *If* $1 \le m \le 2+p$ *and* m *is odd, then* $\beta_k = C^*\alpha_k$ *and* w_k *satisfies*

$$\begin{aligned} w_k :=& \big[\psi_1 + C^*\psi_2\big]\alpha_k + \phi\,\alpha_k\varepsilon_k^m \\ &+\big[\psi_3 + (C^*)^2\psi_4 + C^*\psi_5\big]\alpha_k^2 + o(\alpha_k\varepsilon_k^m) \ \ as\ \ k\to\infty, \end{aligned} \tag{4.16}$$

where the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ *is as in (3.38).*

3. *If* $1 \le m < 2+p$ *and* m *is even, consider*

$$\mathcal{S} = \sum_{|\alpha|=m}\int_{\mathbb{R}^2}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w^2. \tag{4.17}$$

Then for the case where $\mathcal{S}=0$*, we have* $\beta_k = C^*\alpha_k$ *and* w_k *satisfies (4.16), where the constant* $C^* \in \mathbb{R}^+\cup\mathbb{R}^-$ *is as in (3.38). However, for the case where* $\mathcal{S}\neq 0$*, we have*

$$\beta_k = C_1^*\varepsilon_k^m, \tag{4.18}$$

and w_k *satisfies*

$$w_k := C_1^*\psi_2\varepsilon_k^m + \psi_1\alpha_k + (C_1^*)^2\psi_4\varepsilon_k^{2m} + o(\varepsilon_k^{\min\{2+p,2m\}}) \ \ as\ \ k\to\infty, \tag{4.19}$$

where the constant $C_1^* \in \mathbb{R}^+\cup\mathbb{R}^-$ *is given by*

$$C_1^* = -\frac{m+p}{(2+p)\lambda^{2+p+m}}\sum_{|\alpha|=m}\int_{\mathbb{R}^2}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w^2. \tag{4.20}$$

4. *If* $m = 2+p$ *is even, then*

$$\beta_k = C_2^*\alpha_k, \tag{4.21}$$

and w_k *satisfies*

$$w_k := \big[\psi_1 + C_2^*\psi_2\big]\alpha_k + \big[\psi_3 + (C_2^*)^2\psi_4 + C_2^*\psi_5 + \phi\big]\alpha_k^2 + o(\alpha_k^2) \ \ as\ \ k\to\infty, \tag{4.22}$$

where the constant $C_2^* \in \mathbb{R}^+\cup\mathbb{R}^-$ *satisfies*

$$C_2^* = \frac{2}{2+p}\Big[2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2 + 2\int_{\mathbb{R}^2} w\phi\Big]. \tag{4.23}$$

Proof. The same argument of proving Lemma 3.4 yields that

$$\int_{\mathbb{R}^2} w\psi_1 = 0, \ \ \int_{\mathbb{R}^2} w\psi_2 = 0 \ \text{ and } \ I = \int_{\mathbb{R}^2}\big(2w\psi_4 + \psi_2^2\big) = 0, \tag{4.24}$$

and

$$II = 2\int_{\mathbb{R}^2} w\psi_5 + 2\int_{\mathbb{R}^2}\psi_1\psi_2 = -\frac{2+p}{2} < 0. \tag{4.25}$$

It thus follows from (3.40) and Lemma 4.2 that

$$
\begin{aligned}
0 &= 2\int_{\mathbb{R}^2} ww_k + \int_{\mathbb{R}^2} w_k^2 \\
&= 2\int_{\mathbb{R}^2} w\big(\alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 + \alpha_k\varepsilon_k^m\phi + \alpha_k\beta_k\psi_5\big) \\
&\quad + \int_{\mathbb{R}^2} \big(\alpha_k\psi_1 + \beta_k\psi_2 + \alpha_k^2\psi_3 + \beta_k^2\psi_4 + \alpha_k\varepsilon_k^m\phi + \alpha_k\beta_k\psi_5\big)^2 \\
&\quad + o\big([\alpha_k+\beta_k]^2 + \alpha_k\varepsilon_k^m\big) \\
&= \alpha_k\Big(2\int_{\mathbb{R}^2} w\psi_1\Big) + \beta_k\Big(2\int_{\mathbb{R}^2} w\psi_2\Big) + \beta_k^2\Big(2\int_{\mathbb{R}^2} w\psi_4 + \int_{\mathbb{R}^2}\psi_2^2\Big) \\
&\quad + \alpha_k\beta_k\Big(2\int_{\mathbb{R}^2} w\psi_5 + 2\int_{\mathbb{R}^2}\psi_1\psi_2\Big) + \alpha_k^2\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) \\
&\quad + \alpha_k\varepsilon_k^m\Big(2\int_{\mathbb{R}^2} w\phi\Big) + o\big([\alpha_k+\beta_k]^2 + \alpha_k\varepsilon_k^m\big) \\
&= -\frac{2+p}{2}\alpha_k\beta_k + \alpha_k^2\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) + \alpha_k\varepsilon_k^m\Big(2\int_{\mathbb{R}^2} w\phi\Big) \\
&\quad + o\big([\alpha_k+\beta_k]^2 + \alpha_k\varepsilon_k^m\big),
\end{aligned}
\tag{4.26}
$$

where (4.24) and (4.25) are used in the last equality. Following (4.26), we next carry out the proof by considering separately the following four cases:

Case 1. $m > 2+p$. In this case, it follows from (4.26) that

$$
2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2 \neq 0, \tag{4.27}
$$

and moreover,

$$
-\frac{2+p}{2}\beta_k + \alpha_k\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2\Big) = 0, \quad i.e., \ \beta_k = C^*\alpha_k,
$$

where the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is as in (3.38). Finally, the expansion (4.15) follows directly from (4.14) and Lemma 4.2, and Case 1 is therefore proved.

Case 2. $1 \leq m \leq 2+p$ *and* m *is odd.* In this case, since m is odd and $h(x)$ is symmetric about the origin, we obtain from (3.10) and (4.10) that

$$
\begin{aligned}
2\int_{\mathbb{R}^2} w\phi &= 2\int_{\mathbb{R}^2} \phi\mathcal{L}\psi_2 = 2\int_{\mathbb{R}^2} \psi_2\mathcal{L}\phi \\
&= \frac{1}{\lambda^{2+p}}\int_{\mathbb{R}^2}\Big\{\big[x_0\cdot\nabla h(x)\big]g(0)w \\
&\quad + \frac{1}{\lambda^m}\sum_{|\alpha|=m}\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big]h(x)w\Big\}\big(w + x\cdot\nabla w\big) = 0.
\end{aligned}
$$

We then derive from (4.26) that (4.27) still holds and thus $\beta_k = C^*\alpha_k$, where the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is given again by (3.38). Further, the expansion (4.16) follows directly from (4.7) and (4.14).

Case 3. $1 \le m < 2+p$ *and* m *is even.* Since m is even, then $x_0 = 0$ holds in (4.10). Further, since $x^\alpha h(x)$ is homogeneous of degree $m+p$, we then obtain from (4.10) that

$$\begin{aligned}
2\int_{\mathbb{R}^2} w\phi &= 2\int_{\mathbb{R}^2} \phi\mathcal{L}\psi_2 = 2\int_{\mathbb{R}^2} \psi_2\mathcal{L}\phi \\
&= \frac{1}{\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} \Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x) w\big(w + x\cdot\nabla w\big) \\
&= \frac{1}{\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} \Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x) w^2 \\
&\quad + \frac{1}{2\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} \Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x) x\cdot\nabla w^2 \\
&= \frac{1}{\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} \Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x) w^2 \\
&\quad - \frac{1}{2\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} w^2 \Big\{ 2\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0) h(x)\Big] \\
&\quad + x\cdot\nabla\Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0) h(x)\Big]\Big\} \\
&= -\frac{m+p}{2\lambda^{2+p+m}} \sum_{|\alpha|=m} \int_{\mathbb{R}^2} \Big[\frac{x^\alpha}{\alpha!}\mathcal{D}^\alpha g(0)\Big] h(x) w^2 := -\frac{m+p}{2\lambda^{2+p+m}}\mathcal{S},
\end{aligned} \tag{4.28}$$

where $\mathcal{S}$ is as in (4.17). Therefore, if $\mathcal{S} = 0$, then we are in the same situation as that of above Case 2, which gives that $\beta_k = C^*\alpha_k$ and w_k satisfies (4.16), where the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ satisfies (3.38).

We next consider the case where $\mathcal{S} \neq 0$. By applying (4.28), in this case we derive from (4.26) that

$$-\frac{2+p}{2}\alpha_k\beta_k + \alpha_k\varepsilon_k^m\Big(2\int_{\mathbb{R}^2} w\phi\Big) = 0,$$

which implies that $\beta_k = C_1^*\varepsilon_k^m$, where the constant $C_1^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ satisfies (4.20) in view of (4.28). Further, the expansion (4.19) follows directly from (4.18) and Lemma 4.2.

Case 4. $m = 2+p$ *is even.* In this case, we derive from (4.26) that

$$-\frac{2+p}{2}\alpha_k\beta_k + \alpha_k^2\Big(2\int_{\mathbb{R}^2} w\psi_3 + \int_{\mathbb{R}^2}\psi_1^2 + 2\int_{\mathbb{R}^2} w\phi\Big) = 0,$$

which gives that $\beta_k = C_2^*\alpha_k$, where the constant $C_2^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ satisfies (4.23). Further, the expansion (4.22) follows directly from (4.21) and Lemma 4.2. □

Applying directly Lemmas 4.1 and 4.2 as well as Proposition 4.3, we now conclude the following refined spike profiles. Recall that $\lambda > 0$ is defined by (2.2), $\psi_1(x), \cdots, \psi_5(x)$ are given in Lemmas 3.1 and 4.2, and ϕ is given by (4.10).

Theorem 4.4. *Suppose* $V(x) = g(x)h(x)$ *satisfies the assumption* (V) *for* $p > 0$ *and* $m \in \mathbb{N} \cup \{+\infty\}$. *Let* u_a *be a positive minimizer of (1.1) for* $a < a^*$. *Then for any sequence* $\{a_k\}$ *with* $a_k \nearrow a^*$ *as* $k \to \infty$, *there exist a subsequence, still denoted by* $\{a_k\}$, *of* $\{a_k\}$ *and* $\{x_k\} \subset \mathbb{R}^2$ *such that the subsequence* $\{u_k\}$ *satisfies for* $\varepsilon_k := (a^* - a_k)^{\frac{1}{2+p}}$,

1. *If $m > 2+p$, then we have*

$$u_k(x) = \frac{\lambda}{\|w\|_2}\Big\{\frac{1}{\varepsilon_k} w\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+p}\Big[\psi_1 + C^*\psi_2\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{3+2p}\Big[\psi_3 + (C^*)^2\psi_4 + C^*\psi_5\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big)\Big\} + o(\varepsilon_k^{3+2p}) \ \ as \ \ k\to\infty \tag{4.29}$$

uniformly in $\mathbb{R}^2$, where $\{x_k\}$ satisfies

$$\frac{x_k}{\varepsilon_k} = O(\varepsilon_k^m y_0) + o(\varepsilon_k^{2+p}) \ \ as \ \ k\to\infty \tag{4.30}$$

for some $y_0 \in \mathbb{R}^2$, and the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is given by (3.38). Further, if m is even, then $\{x_k\}$ satisfies

$$\frac{x_k}{\varepsilon_k^{3+p}} = o(1) \ \ as \ \ k\to\infty. \tag{4.31}$$

2. *If $1 \le m \le 2+p$ and m is odd, then we have*

$$u_k(x) = \frac{\lambda}{\|w\|_2}\Big\{\frac{1}{\varepsilon_k} w\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+p}\Big[\psi_1 + C^*\psi_2\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{3+2p}\big[\psi_3 + (C^*)^2\psi_4 + C^*\psi_5\big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+m+p}\phi\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big)\Big\} + o(\varepsilon_k^{1+m+p}) \ \ as \ \ k\to\infty \tag{4.32}$$

uniformly in $\mathbb{R}^2$, where $\{x_k\}$ satisfies

$$\frac{x_k}{\varepsilon_k^{m+1}} = O(y_0) \ \ as \ \ k\to\infty. \tag{4.33}$$

for some $y_0 \in \mathbb{R}^2$, and the constant $C^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is given by (3.38).

3. *If $m = 2+p$ is even, then we have*

$$u_k(x) = \frac{\lambda}{\|w\|_2}\Big\{\frac{1}{\varepsilon_k} w\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+p}\Big[\psi_1 + C_2^*\psi_2\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{3+2p}\Big[\psi_3 + (C_2^*)^2\psi_4 + C_2^*\psi_5 + \phi\Big]\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big)\Big\} + o(\varepsilon_k^{3+2p}) \ \ as \ \ k\to\infty \tag{4.34}$$

uniformly in $\mathbb{R}^2$, where $\{x_k\}$ satisfies (4.31) and the constant $C_2^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is defined by (4.23).

4. *If $1 \le m < 2+p$ and m is even, let the constant $\mathcal{S}$ be defined in (4.17). Then for the case where $\mathcal{S} = 0$, u_k satisfies (4.32) and $\{x_k\}$ satisfies*

$$\frac{x_k}{\varepsilon_k^{m+1}} = o(1) \ \ as \ \ k\to\infty. \tag{4.35}$$

However, for the case where $\mathcal{S} \neq 0$, u_k satisfies

$$\begin{aligned} u_k(x) = \frac{\lambda}{\|w\|_2}\Big\{ & \frac{1}{\varepsilon_k} w\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{m-1} C_1^* \psi_2\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) \\ & + \varepsilon_k^{2m-1} (C_1^*)^2 \psi_4\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big) + \varepsilon_k^{1+p} \psi_1\Big(\frac{\lambda(x-x_k)}{\varepsilon_k}\Big)\Big\} \\ & + o(\varepsilon_k^{\min\{2+p,2m\}-1}) \ \ as \ \ k \to \infty \end{aligned} \tag{4.36}$$

uniformly in $\mathbb{R}^2$, where $\{x_k\}$ satisfies (4.35), and the constant $C_1^* \in \mathbb{R}^+ \cup \mathbb{R}^-$ is defined by (4.20).

Proof. (1). If $m > 2+p$, then (4.29) follows directly from Proposition 4.3(1), and (4.30) follows from Lemma 4.1. Specially, if m is even, then Lemma 4.1 gives $y_0 = 0$, and therefore (4.30) implies (4.31).

(2). If $1 \leq m \leq 2+p$ and m is odd, then Proposition 4.3(2) gives (4.32). Moreover, it yields from (4.2) that x_k satisfies $\frac{x_k}{\varepsilon_k} = O(\varepsilon_k^m y_0) + o(\varepsilon_k^m)$ as $k \to \infty$, which then implies (4.33) for some $y_0 \in \mathbb{R}^2$.

(3). If $m = 2+p$ is even, then Proposition 4.3(4) gives (4.34), and we reduce from (4.1) that x_k satisfies (4.31).

(4). If $1 \leq m < 2+p$ and m is even, it then follows from (4.1) that x_k always satisfies (4.35). Moreover, Proposition 4.3(3) gives that if $\mathcal{S} = 0$, then u_k satisfies (4.32); if $\mathcal{S} \neq 0$, then u_k satisfies (4.36). □

We finally remark that Theorem 3.6 and Theorem 4.4(1) with $m = +\infty$ give directly the estimates (1.13) and (1.14) of Theorem 1.1.

A Appendix: The Proof of Lemma 3.4

In this appendix, we shall follow Lemmas 3.1 and 3.3 to address the proof of Lemma 3.4, i.e., (3.34)–(3.36).

The proof of (3.34). Under the assumptions of Lemma 3.4, we first note that the equation (3.7) can be simplified as

$$\nabla\psi_1(0) = 0, \quad \mathcal{L}\psi_1 = -\frac{2w^3}{\int_{\mathbb{R}^2} w^4} - \frac{2h(x)w}{p\int_{\mathbb{R}^2} h(x)w^2} \ \text{ in } \mathbb{R}^2, \tag{A.1}$$

due to the fact that

$$a^* = \|w\|_2^2 = \frac{1}{2}\int_{\mathbb{R}^2} w^4. \tag{A.2}$$

By (3.10) and (A.1), we then have

$$\begin{aligned}
2\int_{\mathbb{R}^2} w\psi_1 &= 2\int_{\mathbb{R}^2} \mathcal{L}\psi_2\psi_1 = 2\int_{\mathbb{R}^2} \psi_2\mathcal{L}\psi_1 \\
&= \int_{\mathbb{R}^2} \Big[\frac{2w^3}{\int_{\mathbb{R}^2} w^4} + \frac{2h(x)w}{p\int_{\mathbb{R}^2} h(x)w^2}\Big]\big(w + x\cdot\nabla w\big) \\
&= 2 + \frac{2}{p} + \frac{2}{\int_{\mathbb{R}^2} w^4}\int_{\mathbb{R}^2} w^3\big(x\cdot\nabla w\big) + \frac{2}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2} h(x)w\big(x\cdot\nabla w\big) \\
&= 2 + \frac{2}{p} + \frac{1}{2\int_{\mathbb{R}^2} w^4}\int_{\mathbb{R}^2} \big(x\cdot\nabla w^4\big) + \frac{1}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2} h(x)\big(x\cdot\nabla w^2\big) \\
&= 2 + \frac{2}{p} - 1 - \frac{1}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2} w^2\Big[2h(x) + \big(x\cdot\nabla h(x)\big)\Big] \\
&= 2 + \frac{2}{p} - 1 - \frac{(p+2)}{p} = 0,
\end{aligned}$$

since $x\cdot\nabla h(x) = ph(x)$ holds in $\mathbb{R}^2$. Also, we deduce from (3.10) that

$$2\int_{\mathbb{R}^2} w\psi_2 = -\int_{\mathbb{R}^2} w(w + x\cdot\nabla w) = -\int_{\mathbb{R}^2} w^2 - \frac{1}{2}\int_{\mathbb{R}^2} (x\cdot\nabla w^2) = 0,$$

which thus completes the proof of (3.34). □

The proof of (3.35). By Lemmas 3.1 and 3.3, we obtain that

$$\begin{aligned}
I &= \int_{\mathbb{R}^2} \big(2w\psi_4 + \psi_2^2\big) = \int_{\mathbb{R}^2} \psi_2^2 + 2\langle\mathcal{L}\psi_2, \psi_4\rangle \\
&= \int_{\mathbb{R}^2} \psi_2^2 + 2\langle\psi_2, \mathcal{L}\psi_4\rangle = \int_{\mathbb{R}^2} \psi_2^2 + 2\langle\psi_2, (3w\psi_2^2 + \psi_2)\rangle \\
&= 3\int_{\mathbb{R}^2} \psi_2^2 + 6\int_{\mathbb{R}^2} w\psi_2^3,
\end{aligned}$$

which implies that

$$\begin{aligned}
\frac{4I}{3} \div 2\pi &= 4\Big[\int_{\mathbb{R}^2} \psi_2^2 + 2\int_{\mathbb{R}^2} w\psi_2^3\Big] \div 2\pi \\
&= \int_0^\infty r(w + rw')^2 - \int_0^\infty rw(w + rw')^3 := A - B.
\end{aligned} \tag{A.3}$$

Here we have

$$\begin{aligned}
A = \int_0^\infty r(w + rw')^2 &= \int_0^\infty r^3(w')^2 + \int_0^\infty rw^2 + \int_0^\infty r^2 dw^2 \\
&= \int_0^\infty r^3(w')^2 - \frac{1}{2}\int_0^\infty rw^4,
\end{aligned}$$

where (A.2) is used, and

$$\begin{aligned}
B &= \int_0^\infty rw(w + rw')^3 \\
&= \Big[\int_0^\infty rw^4 + 3\int_0^\infty r^2w^3w'\Big] + 3\int_0^\infty r^3w^2w'w' + \int_0^\infty r^4w(w')^3 \\
&= -\frac{1}{2}\int_0^\infty rw^4 + 3\int_0^\infty r^3w^2w'w' + \int_0^\infty r^4w(w')^3.
\end{aligned}$$

Therefore, we get from (A.3) that

$$\frac{4I}{3} \div 2\pi = \int_0^\infty r^3 (w')^2 - \int_0^\infty r^4 w (w')^3 - 3\int_0^\infty r^3 w^2 w' w' := C + D + E. \tag{A.4}$$

To further simplify I, recall that

$$rw'' = -w' + rw - rw^3, \tag{A.5}$$

by which we then have

$$\begin{aligned} C = \int_0^\infty r^3 w' dw &= -\int_0^\infty w(r^3 w')' \\ &= -\int_0^\infty w\big[3r^2 w' + r^2(-w' + rw - rw^3)\big] \\ &= -\int_0^\infty w\big[2r^2 w' + r^3 w - r^3 w^3\big] \\ &= 2\int_0^\infty rw^2 - \int_0^\infty r^3 w^2 + \int_0^\infty r^3 w^4. \end{aligned}$$

Similarly, we have

$$\begin{aligned} D = -\frac{1}{2}\int_0^\infty r^4 (w')^2 dw^2 &= \frac{1}{2}\int_0^\infty w^2\big[4r^3 (w')^2 + 2r^3 w'(-w' + rw - rw^3)\big] \\ &= \int_0^\infty r^3 w^2 (w')^2 + \frac{1}{4}\int_0^\infty r^4 dw^4 - \frac{1}{6}\int_0^\infty r^4 dw^6 \\ &= \int_0^\infty r^3 w^2 (w')^2 - \int_0^\infty r^3 w^4 + \frac{2}{3}\int_0^\infty r^3 w^6. \end{aligned}$$

Note from (A.5) that

$$\begin{aligned} -2\int_0^\infty r^3 w^2 (w')^2 = -\frac{2}{3}\int_0^\infty r^3 w' dw^3 &= \frac{2}{3}\int_0^\infty w^3\big[3r^2 w' + r^2(-w' + rw - rw^3)\big] \\ &= \frac{4}{3}\int_0^\infty w^3 r^2 w' + \frac{2}{3}\int_0^\infty r^3 w^4 - \frac{2}{3}\int_0^\infty r^3 w^6 \\ &= -\frac{2}{3}\int_0^\infty rw^4 + \frac{2}{3}\int_0^\infty r^3 w^4 - \frac{2}{3}\int_0^\infty r^3 w^6. \end{aligned}$$

We thus derive that

$$\begin{aligned} D + E &= -2\int_0^\infty r^3 w^2 (w')^2 - \int_0^\infty r^3 w^4 + \frac{2}{3}\int_0^\infty r^3 w^6 \\ &= -\frac{2}{3}\int_0^\infty rw^4 - \frac{1}{3}\int_0^\infty r^3 w^4, \end{aligned}$$

by which we conclude from (A.2) and (A.4) that

$$\frac{4I}{3} \div 2\pi = C + D + E = \frac{1}{3}\Big[2\int_0^\infty rw^2 - 3\int_0^\infty r^3 w^2 + 2\int_0^\infty r^3 w^4\Big]. \tag{A.6}$$

In the following, we note that w satisfies

$$(rw')' = rw - rw^3, \ \ r > 0. \tag{A.7}$$

Multiplying (A.7) by $r^3 w'$ and integrating on $[0, \infty)$, we get that

$$\begin{aligned}\int_0^\infty r^3 w'(rw')' = \int_0^\infty r^3 w'[rw - rw^3] &= \frac{1}{2}\int_0^\infty r^4 dw^2 - \frac{1}{4}\int_0^\infty r^4 dw^4 \\ &= -2\int_0^\infty r^3 w^2 + \int_0^\infty r^3 w^4.\end{aligned}$$

Note also that

$$\int_0^\infty r^3 w'(rw')' = \int_0^\infty r^3 (w')^2 + \frac{1}{2}\int_0^\infty r^4 d(w')^2 = -\int_0^\infty r^3 (w')^2.$$

By combining above two identities, it yields that

$$\int_0^\infty r^3 (w')^2 = 2\int_0^\infty r^3 w^2 - \int_0^\infty r^3 w^4. \tag{A.8}$$

On the other hand, multiplying (A.7) by $r^2 w$ and integrating on $[0, \infty)$, we obtain that

$$\begin{aligned}\int_0^\infty r^3 w^2 - \int_0^\infty r^3 w^4 = \int_0^\infty r^2 w(rw')' &= \int_0^\infty r^2 ww' + \int_0^\infty r^3 w dw' \\ &= \int_0^\infty r^2 ww' - \int_0^\infty w'(3r^2 w + r^3 w') \\ &= -2\int_0^\infty r^2 ww' - \int_0^\infty r^3 (w')^2 \\ &= 2\int_0^\infty rw^2 - \int_0^\infty r^3 (w')^2,\end{aligned}$$

which then implies that

$$\int_0^\infty r^3 (w')^2 = 2\int_0^\infty rw^2 - \int_0^\infty r^3 w^2 + \int_0^\infty r^3 w^4. \tag{A.9}$$

We thus conclude from (A.8) and (A.9) that

$$2\int_0^\infty rw^2 - 3\int_0^\infty r^3 w^2 + 2\int_0^\infty r^3 w^4 = 0,$$

which therefore implies that $I = 0$ in view of (A.6), i.e., (3.35) holds. □

The proof of (3.36). Following Lemmas 3.1 and 3.3 again, we get that

$$\begin{aligned}II &= 2\int_{\mathbb{R}^2} \psi_5 \mathcal{L}\psi_2 + 2\int_{\mathbb{R}^2} \psi_1\psi_2 = 2\int_{\mathbb{R}^2} \psi_2[\mathcal{L}\psi_5 + \psi_1] \\ &= -\int_{\mathbb{R}^2} (w + x\cdot\nabla w)(6w\psi_1\psi_2 + 2\psi_1) \\ &\quad -\frac{1}{2}\int_{\mathbb{R}^2} \Big[\frac{3w^2}{a^*} + \frac{h(x)}{\lambda^{2+p}}\Big]\big(w + x\cdot\nabla w\big)^2 := A + B.\end{aligned} \tag{A.10}$$

Since $x \cdot \nabla h(x) = ph(x)$ holds in $\mathbb{R}^2$, we derive from (A.1) that

$$\begin{aligned}
B &= -\frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{3w^2}{a^*} + \frac{h(x)}{\lambda^{2+p}}\Big]\big[w^2 + 2w(x\cdot\nabla w) + (x\cdot\nabla w)^2\big] \\
&= -\int_{\mathbb{R}^2}\Big[\frac{3w^2}{\int_{\mathbb{R}^2} w^4} + \frac{h(x)}{p\int_{\mathbb{R}^2} h(x)w^2}\Big]\big[w^2 + 2w(x\cdot\nabla w)\big] \\
&\quad -\frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{3w^2}{a^*} + \frac{h(x)}{\lambda^{2+p}}\Big]\big(x\cdot\nabla w\big)^2 \\
&= -3 - \frac{1}{p} - \frac{3}{2\int_{\mathbb{R}^2} w^4}\int_{\mathbb{R}^2}(x\cdot\nabla w^4) - \frac{1}{p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2}\big(h(x)x\cdot\nabla w^2\big) \\
&\quad -\frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{3w^2}{a^*} + \frac{h(x)}{\lambda^{2+p}}\Big]\big(x\cdot\nabla w\big)^2 \\
&= -3 - \frac{1}{p} + 3 + \frac{2+p}{p} + C_0 = \frac{p+1}{p} + C_0,
\end{aligned}$$

where the term C_0 satisfies

$$\begin{aligned}
C_0 &= -\frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{3w^2}{a^*} + \frac{h(x)}{\lambda^{2+p}}\Big]\big(x\cdot\nabla w\big)^2 \\
&= -\frac{1}{2a^*}\int_{\mathbb{R}^2}(x\cdot\nabla w)(x\cdot\nabla w^3) - \frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} h(x)\big(x\cdot\nabla w\big)(x\cdot\nabla w) \\
&= \frac{1}{2a^*}\int_{\mathbb{R}^2} w^3\Big[2(x\cdot\nabla w) + x\cdot\nabla(x\cdot\nabla w)\Big] \\
&\quad +\frac{1}{2\lambda^{2+p}}\int_{\mathbb{R}^2} w\Big\{(2+p)h(x)(x\cdot\nabla w) + h(x)\Big[x\cdot\nabla(x\cdot\nabla w)\Big]\Big\} \\
&= \frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{w^3}{a^*} + \frac{wh(x)}{\lambda^{2+p}}\Big]\big[x\cdot\nabla\big(x\cdot\nabla w\big)\big] \\
&\quad +\frac{1}{a^*}\int_{\mathbb{R}^2} w^3(x\cdot\nabla w) + \frac{2+p}{2\lambda^{2+p}}\int_{\mathbb{R}^2} wh(x)(x\cdot\nabla w) \\
&= \frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{w^3}{a^*} + \frac{wh(x)}{\lambda^{2+p}}\Big]\big[x\cdot\nabla\big(x\cdot\nabla w\big)\big] \\
&\quad +\frac{1}{2\int_{\mathbb{R}^2} w^4}\int_{\mathbb{R}^2}(x\cdot\nabla w^4) + \frac{2+p}{2p\int_{\mathbb{R}^2} h(x)w^2}\int_{\mathbb{R}^2}\big(h(x)x\cdot\nabla w^2\big) \\
&= \frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{w^3}{a^*} + \frac{wh(x)}{\lambda^{2+p}}\Big]\big[x\cdot\nabla\big(x\cdot\nabla w\big)\big] - 1 - \frac{(2+p)^2}{2p},
\end{aligned}$$

in view of (A.1). We thus have

$$B = \frac{1}{2}\int_{\mathbb{R}^2}\Big[\frac{w^3}{a^*} + \frac{wh(x)}{\lambda^{2+p}}\Big]\big[x\cdot\nabla\big(x\cdot\nabla w\big)\big] - \frac{p^2+4p+2}{2p}. \tag{A.11}$$

We next calculate the term A as follows. Observe that

$$\begin{aligned}
&\frac{1}{2}\int_{\mathbb{R}^2}\psi_1 x\cdot\nabla\big(x\cdot\nabla w\big)\\
&=-\frac{1}{2}\int_{\mathbb{R}^2}\big(x\cdot\nabla w\big)\big[2\psi_1+\big(x\cdot\nabla\psi_1\big)\big]\\
&=-\int_{\mathbb{R}^2}\psi_1\big(x\cdot\nabla w\big)-\frac{1}{2}\int_{\mathbb{R}^2}\big(x\cdot\nabla\psi_1\big)\big(x\cdot\nabla w\big)\\
&=-\int_{\mathbb{R}^2}\psi_1\big(x\cdot\nabla w\big)+\frac{1}{2}\int_{\mathbb{R}^2}w\big[2\big(x\cdot\nabla\psi_1\big)+x\cdot\nabla\big(x\cdot\nabla\psi_1\big)\big]\\
&=-\int_{\mathbb{R}^2}\psi_1\big(x\cdot\nabla w\big)+\int_{\mathbb{R}^2}w\big(x\cdot\nabla\psi_1\big)+\frac{1}{2}\int_{\mathbb{R}^2}wx\cdot\nabla\big(x\cdot\nabla\psi_1\big),
\end{aligned}$$

which implies that

$$\begin{aligned}
&-\int_{\mathbb{R}^2}\psi_1\big(x\cdot\nabla w\big)+\int_{\mathbb{R}^2}w\big(x\cdot\nabla\psi_1\big)\\
&=\frac{1}{2}\int_{\mathbb{R}^2}\psi_1 x\cdot\nabla\big(x\cdot\nabla w\big)-\frac{1}{2}\int_{\mathbb{R}^2}wx\cdot\nabla\big(x\cdot\nabla\psi_1\big).
\end{aligned}\tag{A.12}$$

Using (A.12), we then derive that

$$\begin{aligned}
A&=-\int_{\mathbb{R}^2}(w+x\cdot\nabla w)(6w\psi_1\psi_2+2\psi_1)\\
&=-2\int_{\mathbb{R}^2}w\psi_1-2\int_{\mathbb{R}^2}\psi_1(x\cdot\nabla w)+3\int_{\mathbb{R}^2}w\psi_1(w+x\cdot\nabla w)^2\\
&=-2\int_{\mathbb{R}^2}w\psi_1-\int_{\mathbb{R}^2}\psi_1(x\cdot\nabla w)\\
&\quad+\int_{\mathbb{R}^2}w\big[2\psi_1+x\cdot\nabla\psi_1\big]+3\int_{\mathbb{R}^2}w\psi_1(w+x\cdot\nabla w)^2\\
&=-\int_{\mathbb{R}^2}\psi_1(x\cdot\nabla w)+\int_{\mathbb{R}^2}w(x\cdot\nabla\psi_1)+D\\
&=\frac{1}{2}\int_{\mathbb{R}^2}\psi_1 x\cdot\nabla\big(x\cdot\nabla w\big)-\frac{1}{2}\int_{\mathbb{R}^2}wx\cdot\nabla\big(x\cdot\nabla\psi_1\big)+D,
\end{aligned}\tag{A.13}$$

where the term D satisfies

$$\begin{aligned}
D&=3\int_{\mathbb{R}^2}w\psi_1\Big[w^2+2w(x\cdot\nabla w)+(x\cdot\nabla w)^2\Big]\\
&=3\int_{\mathbb{R}^2}w^3\psi_1+6\int_{\mathbb{R}^2}w^2\psi_1(x\cdot\nabla w)+\frac{3}{2}\int_{\mathbb{R}^2}\psi_1(x\cdot\nabla w)(x\cdot\nabla w^2)\\
&=3\int_{\mathbb{R}^2}w^3\psi_1+6\int_{\mathbb{R}^2}w^2\psi_1(x\cdot\nabla w)\\
&\quad-\frac{3}{2}\int_{\mathbb{R}^2}w^2\Big\{2\psi_1(x\cdot\nabla w)+(x\cdot\nabla w)(x\cdot\nabla\psi_1)+\psi_1\Big[x\cdot\nabla(x\cdot\nabla w)\Big]\Big\}.
\end{aligned}$$

Since

$$\begin{aligned}
&-\frac{3}{2}\int_{\mathbb{R}^2} w^2(x\cdot\nabla w)(x\cdot\nabla\psi_1)\\
&=-\frac{1}{2}\int_{\mathbb{R}^2}(x\cdot\nabla\psi_1)(x\cdot\nabla w^3)\\
&=\frac{1}{2}\int_{\mathbb{R}^2} w^3\Big[x\cdot\nabla(x\cdot\nabla\psi_1)+2(x\cdot\nabla\psi_1)\Big]\\
&=\frac{1}{2}\int_{\mathbb{R}^2} w^3 x\cdot\nabla(x\cdot\nabla\psi_1)+\int_{\mathbb{R}^2} w^3(x\cdot\nabla\psi_1)\\
&=\frac{1}{2}\int_{\mathbb{R}^2} w^3 x\cdot\nabla(x\cdot\nabla\psi_1)-\int_{\mathbb{R}^2}\psi_1\Big[2w^3+3w^2(x\cdot\nabla w)\Big]\\
&=\frac{1}{2}\int_{\mathbb{R}^2} w^3 x\cdot\nabla(x\cdot\nabla\psi_1)-2\int_{\mathbb{R}^2} w^3\psi_1-3\int_{\mathbb{R}^2} w^2\psi_1(x\cdot\nabla w),
\end{aligned}$$

the term D can be further simplified as

$$D=\int_{\mathbb{R}^2} w^3\psi_1-\frac{3}{2}\int_{\mathbb{R}^2} w^2\psi_1\big[x\cdot\nabla(x\cdot\nabla w)\big]+\frac{1}{2}\int_{\mathbb{R}^2} w^3 x\cdot\nabla(x\cdot\nabla\psi_1). \tag{A.14}$$

Applying (A.14), we then obtain from (A.13) that

$$\begin{aligned}
A=&\int_{\mathbb{R}^2} w^3\psi_1+\frac{1}{2}\int_{\mathbb{R}^2}(1-3w^2)\psi_1\big[x\cdot\nabla(x\cdot\nabla w)\big]\\
&-\frac{1}{2}\int_{\mathbb{R}^2}\Delta w\big[x\cdot\nabla(x\cdot\nabla\psi_1)\big],
\end{aligned} \tag{A.15}$$

since w solves the equation $w^3-w=-\Delta w$ in $\mathbb{R}^2$.

Combining (A.11) and (A.15) now yields that

$$\begin{aligned}
II=A+B=&\int_{\mathbb{R}^2} w^3\psi_1-\frac{p^2+4p+2}{2p}\\
&+\frac{1}{2}\int_{\mathbb{R}^2}\big[x\cdot\nabla(x\cdot\nabla w)\big]\Delta\psi_1-\frac{1}{2}\int_{\mathbb{R}^2}\big[x\cdot\nabla(x\cdot\nabla\psi_1)\big]\Delta w.
\end{aligned} \tag{A.16}$$

We claim that

$$\int_{\mathbb{R}^2} w^3\psi_1=\frac{p+1}{p}. \tag{A.17}$$

Actually, multiplying (A.1) by w and integrating on $\mathbb{R}^2$ gives that

$$\int_{\mathbb{R}^2}\nabla\psi_1\nabla w-3\int_{\mathbb{R}^2} w^3\psi_1=-\int_{\mathbb{R}^2}\Big[\frac{2w^4}{\int_{\mathbb{R}^2} w^4}+\frac{2h(x)w^2}{p\int_{\mathbb{R}^2} h(x)w^2}\Big]=-\frac{2(p+1)}{p},$$

due to the fact that $\int_{\mathbb{R}^2} w\psi_1=0$ by (3.34). On the other hand, multiplying (1.4) by ψ_1 and integrating on $\mathbb{R}^2$ gives that

$$\int_{\mathbb{R}^2}\nabla\psi_1\nabla w=-\int_{\mathbb{R}^2} w\psi_1+\int_{\mathbb{R}^2} w^3\psi_1=\int_{\mathbb{R}^2} w^3\psi_1.$$

The claim (A.17) then follows directly from above two identities. We next claim that

$$\int_{\mathbb{R}^2}\big[x\cdot\nabla(x\cdot\nabla w)\big]\Delta\psi_1=\int_{\mathbb{R}^2}\big[x\cdot\nabla(x\cdot\nabla\psi_1)\big]\Delta w. \tag{A.18}$$

To prove (A.18), rewrite ψ_1 as $\psi_1(x) = \psi_1(r,\theta)$, where (r,θ) is the polar coordinate in $\mathbb{R}^2$, such that

$$\Delta\psi_1 = \big(\psi_1\big)_{rr} + \frac{1}{r}\big(\psi_1\big)_r + \frac{1}{r^2}\big(\psi_1\big)_{\theta\theta}, \quad \nabla\psi_1 = \frac{x}{r}\big(\psi_1\big)_r + \frac{x^\perp}{r^2}\big(\psi_1\big)_\theta, \tag{A.19}$$

where $x^\perp = (-x_2, x_1)$ for $x = (x_1, x_2) \in \mathbb{R}^2$. We then derive from (3.7) that

$$\begin{aligned}\int_{\mathbb{R}^2} \big[x\cdot\nabla(x\cdot\nabla w)\big]\Delta\psi_1 &= \int_0^{2\pi}\int_0^\infty r(rw')'\Big\{\big[r\big(\psi_1\big)_r\big]_r + \frac{\big(\psi_1\big)_{\theta\theta}}{r}\Big\}drd\theta \\ &= \int_0^{2\pi}\int_0^\infty r(rw')'\big[r\big(\psi_1\big)_r\big]_r drd\theta + \int_0^{2\pi}\int_0^\infty (rw')'\big(\psi_1\big)_{\theta\theta}drd\theta \\ &= \int_0^{2\pi}\int_0^\infty r(rw')'\big[r\big(\psi_1\big)_r\big]_r drd\theta,\end{aligned}$$

and

$$\int_{\mathbb{R}^2} \big[x\cdot\nabla(x\cdot\nabla\psi_1)\big]\Delta w = \int_0^{2\pi}\int_0^\infty r\big[r\big(\psi_1\big)_r\big]_r(rw')'drd\theta,$$

which thus imply that (A.18) holds. Applying (A.17) and (A.18), we therefore conclude from (A.16) that

$$II = \frac{p+1}{p} - \frac{p^2+4p+2}{2p} = -\frac{2+p}{2},$$

which gives (3.36), and the proof is complete. □

Acknowledgements: The first author thanks Prof. Robert Seiringer for fruitful discussions on the present work. Part of the present work was finished when the first author was visiting Taida Institute of Mathematical Sciences (TIMS) in October 2013 and Pacific Institute for Mathematical Sciences (PIMS) from March to April in 2016. He would like to thank both institutes for their warm hospitality.

References

[1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman and E. A. Cornell, *Observation of Bose-Einstein condensation in a dilute atomic vapor*, Science **269** (1995), 198–201.

[2] W. Z. Bao and Y. Y. Cai, *Mathematical theory and numerical methods for Bose-Einstein condensation*, Kinetic and Related Models **6** (2013), 1–135.

[3] D. M. Cao, S. L. Li, and P. Luo, *Uniqueness of positive bound states with multi-bump for nonlinear Schrödinger equations*, Calc. Var. Partial Differential Equations **54** (2015), no. 4, 4037–4063.

[4] T. Cazenave, Semilinear Schrödinger Equations, Courant Lecture Notes in Math. **10**, Courant Institute of Mathematical Science/AMS, New York, (2003).

[5] F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari, *Theory of Bose-Einstein condensation in trapped gases*, Rev. Modern Phys. **71** (1999), 463–512.

[6] Y. B. Deng, C. S. Lin, and S. Yan, *On the prescribed scalar curvature problem in $\mathbb{R}^N$, local uniqueness and periodicity*, J. Math. Pures Appl. **104** (2015), no. 6, 1013–1044.

[7] B. Gidas, W. M. Ni and L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in $\mathbb{R}^n$*, Mathematical analysis and applications Part A, Adv. in Math. Suppl. Stud. vol. **7** (1981), 369–402.

[8] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, (1997).

[9] M. Grossi, *On the number of single-peak solutions of the nonlinear Schrödinger equations*, Ann. Inst H. Poincar Anal. Non Linaire **19** (2002), 261–280.

[10] Y. J. Guo and R. Seiringer, *On the mass concentration for Bose-Einstein condensates with attractive interactions*, Lett. Math. Phys. **104** (2014), 141–156.

[11] Y. J. Guo, Z. Q. Wang, X. Y. Zeng and H. S. Zhou, *Properties for ground states of attractive Gross-Pitaevskii equations with multi-well potentials*, arXiv:1502.01839, submitted, (2015).

[12] Y. J. Guo, X. Y. Zeng and H. S. Zhou, *Energy estimates and symmetry breaking in attractive Bose-Einstein condensates with ring-shaped potentials*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), 809-828.

[13] Y. Kagan, A. E. Muryshev and G. V. Shlyapnikov, *Collapse and Bose-Einstein condensation in a trapped Bose gas with nagative scattering length*, Phys. Rev. Lett. **81** (1998), 933–937.

[14] M. K. Kwong, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in $\mathbb{R}^N$*, Arch. Rational Mech. Anal. **105** (1989), 243–266.

[15] E. H. Lieb and M. Loss, Analysis, Graduate Studies in Math. **14**, Amer. Math. Soc., Providence, RI, second edition (2001).

[16] T. C. Lin and J. C. Wei, *Orbital stability of bound states of semiclassical nonlinear Schrödinger equations with critical nonlinearity*, SIAM J. Math. Anal. **40** (2008), no. 1, 365–381.

[17] T. C. Lin, J. C. Wei and W. Yao, *Orbital stability of bound states of nonlinear Schrödinger equations with linear and nonlinear optical lattices*, J. Differential Equations **249** (2010), no. 9, 2111–2146.

[18] M. Maeda, *On the symmetry of the ground states of nonlinear Schrödinger equation with potential*, Adv. Nonlinear Stud. **10** (2010), 895–925.

[19] K. McLeod and J. Serrin, *Uniqueness of positive radial solutions of $\Delta u + f(u) = 0$ in $\mathbb{R}^n$*, Arch. Rational Mech. Anal. **99** (1987), 115–145.

[20] W.-M. Ni and I. Takagi, *On the shape of least-energy solutions to a semilinear Neumann problem*, Comm. Pure Appl. Math. **44** (1991), 819–851.

[21] W.-M. Ni and J. C. Wei, *On the location and profile of spike-layer solutions to singularly perturbed semilinear Dirichlet problems*, Comm. Pure Appl. Math. **48** (1995), 731–768.

[22] M. Reed and B. Simon, Methods of Modern Mathematical Physics. IV. Analysis of Operators, Academic Press, New York-London, 1978.

[23] C. A. Sackett, H. T. C. Stoof, R. G. Hulet, *Growth and collapse of a Bose-Einstein condensate with attractive interactions*, Phys. Rev. Lett. **80**, 2031 (1998).

[24] X. F. Wang, *On concentration of positive bound states of nonlinear Schrödinger equations*, Comm. Math. Phys. **153** (1993), 229–244.

[25] M. I. Weinstein, *Nonlinear Schrödinger equations and sharp interpolations estimates*, Comm. Math. Phys. **87** (1983), 567–576.

[26] J. Zhang, *Stability of attractive Bose-Einstein condensates*, J. Stat. Phys. **101** (2000), 731–746.